

FUNCTIONALIZED TGF- β FUSION PROTEINS

CROSS REFERENCE TO RELATED APPLICATION

This application claims the benefit of U.S. Provisional Application No.
5 60/242,292, filed October 20, 2000. The provisional application is incorporated
herein in its entirety.

FIELD

The disclosure is in the field of protein labeling and detection. More
10 specifically, the disclosure relates to the use of molecular tags to identify and track
recombinant molecules, such as small proteins, in a subject.

BACKGROUND

TGF- β Protein Family

15 Transforming growth factor- β 1 (TGF- β 1) is a growth factor and
immunomodulatory cytokine that is secreted from cells and acts through specific
binding interactions with a collection of different cell-surface localized receptors.
TGF- β 1 is the prototype for a large family of secreted polypeptides that includes the
three mammalian TGF- β isoforms (TGF- β 1, TGF- β 2, and TGF- β 3), bone
20 morphogenesis proteins (BMPs), activins, and Müllerian inhibitory substance (MIS).
More distantly related members of this protein family include murine *nodal* gene
products, *Drosophila* decapentaplegic complex gene products, and Vg1 from
Xenopus.

In general, TGF- β family proteins are homodimers, wherein each functional
25 protein complex includes two identical, associated monomer subunits. The crystal
structure of the TGF- β 1 homodimer is known (Hinck *et al.*, *Biochem.*, 35:8517-
8534, 1996; Qian *et al.*, *J. Biol. Chem.*, 271:30656-30662, 1996). TGF- β is a very
compact protein, having four intramolecular disulfide bridges within each subunit,
as well as one intermolecular disulfide bridge.

30 Each monomer of the protein is synthesized as a large (~55 kDa) precursor
molecule with a long (about 278 residue) N-terminal pro-region and a much shorter

(112 residue, 12.5 kDa) C-terminal active domain (the mature region). During the maturation process, two precursor molecules associate with each other; the pro-region is important for proper folding of and proper association between the two active domain monomers. The pro-region of each monomer is proteolytically
5 cleaved from the associated active domain; in most instances however, the pro-region remains associated with the mature TGF- β fragment. The severed pro-region is referred to as the "latency-associated peptide" (LAP). LAP is responsible for blocking the correctly folded TGF- β homodimer so that it does not interact with its receptor. For an excellent discussion of TGF- β synthesis, see Khalil, *Micro. Infect.*,
10 1:1255-1263, 1999.

TGF- β s and their receptors are expressed in essentially all tissues, and have been found to be important in many cellular processes. These include cell growth and differentiation, immunosuppression, inflammation, and the expression of extracellular matrix proteins. By way of example, in animal models TGF- β has been
15 shown to attenuate the symptoms associated with various diseases and disorders, including rheumatoid arthritis, multiple sclerosis, wound healing, bronchial asthma, and inflammatory bowel disease, and has been used in the clinical setting to enhance wound healing.

TGF- β 1 was the first identified member of the TGF- β family, and has been
20 intensely studied for over 20 years. There are some TGF- β 1 antibodies available, but their usefulness in a clinical setting is limited at least in part because they often display some degree of cross-reactivity to other TGF- β family proteins (see, *e.g.*, U.S. Patent No. 5,571,714). In most experiments, TGF- β is iodinated with ^{125}I to enable researchers to track the protein. Radioactive iodination is an expensive and
25 hazardous process, and it usually would be inappropriate to use ^{125}I labeled proteins for *in vivo* experimentation, for instance in clinical trials.

The ability to track the distribution of any exogenously administered, recombinant forms of TGF- β family proteins has been restricted by the inability to distinguish between the endogenous forms of the protein produced in treated cells or
30 tissues. In addition, available antibodies to these proteins exhibit some degree of cross-reactivity with related family members.

There have been a few reports of TGF- β fusions in the literature, but the described molecules have been essentially biologically non-functional. In an effort to produce large quantities of easily purified TGF- β that retained activity, Nimni and co-workers expressed 6x His-tagged TGF- β fusion proteins in *Escherichia coli* (Tuan *et al.*, *Conn. Tiss. Res.*, 34:1-9, 1996; Han *et al.*, *Prot. Expr. Purif.*, 11:169-178, 1997). Serious difficulties were encountered in refolding the denatured fusion protein, and full biological activity was not retained using this system. In addition, the Nimni constructs cannot be used to express a tagged TGF- β in a mammalian host, since the constructs lack a part the TGF- β pro-protein (the LAP), which is essential for secretion and proper folding of the TGF- β protein. In an earlier effort, Wakefield *et al.* (*Growth Factors*, 5:243-253, 1991) reported attaching an endoplasmic reticulum retention signal (KDEL) to the C-terminus of full-length TGF- β 1, in an attempt to maintain the protein in the cell (rather than secreting it to the extracellular matrix). This construct had no biological activity.

It is believed that all prior efforts to fuse a TGF- β family protein to a peptide or protein have resulted in biologically non-functional molecules. Therefore a need still exists for TGF- β family protein fusions that maintain substantial biological activity.

SUMMARY OF THE DISCLOSURE

This disclosure provides functionalized TGF- β fusion proteins that maintain substantial TGF- β biological activity. These fusion proteins are achieved by placing a functionalizing peptide between the pro- and active (mature) portions of a TGF- β protein, or at a relatively non-conserved site within the mature region of a TGF- β protein.

Encompassed herein are functional TGF- β family fusion proteins that contain a functionalizing peptide portion for detecting, quantifying or providing a specific additional function to the fusion protein and a mature TGF- β family protein, both as a monomer and in the form of a dimer (*e.g.*, a homodimer). Also encompassed are nucleic acid molecules encoding such fusion proteins, and

conservative substitutions of such molecules. This disclosure also provides methods for making and using the fusion proteins described, as well as kits.

The foregoing and other features and advantages will become more apparent from the following detailed description of several embodiments, which proceeds
5 with reference to the accompanying figures.

BRIEF DESCRIPTION OF THE FIGURES

FIG 1 shows a phylogenetic tree illustrating the relatedness of several members of the TGF- β superfamily of proteins.

10 FIG 2 is a series of schematic drawings depicting (FIG 2A) the structure of a native generic TGF- β protein, and (FIG 2B) the design of two examples of functionalized TGF- β fusion proteins, NFLAG and 11/12FLAG, showing the points of insertion of the FLAG tag.

FIG 3 shows the graphical maps of plasmids encoding the NFLAG and
15 11/12FLAG fusions. The respective plasmids pNFLAGb1 (FIG 3A) and p11/12FLABb1 (FIG 3B) are derived from the pEBB plasmid (Mayer *et al.*, *Curr. Biol.*, 5:296-305, 1995).

FIG 4 shows the quantification and activity of functionalized TGF- β fusion proteins. FIG 4A is a graph showing the amount of the indicated TGF- β protein
20 secreted into COS cell supernatants, as measured in media conditioned by COS cells using a standard ELISA kit specific for TGF- β 1 (R&D Systems, Minneapolis, MN). FIG 4B is a series of Western blots; panels A (anti-TGF- β 1 primary antibody) and B (anti-FLAG primary antibody) show the presence of the indicated TGF- β and TGF- β fusion proteins in mature and precursor forms.

25 Panel C of FIG 4B shows a Western blot (anti-phospho-smad2 primary antibody) demonstrating the induction of phosphorylation of Smad2 with media conditioned by COS cells expressing the indicated TGF- β proteins.

FIG 5 is a bar graph quantifying and comparing the ability of the indicated dilutions of COS cell supernatants (pEBB, WT, NFLAG, 11/12 FLAG,
30 differentiated by the indicated grey tones) to reversibly inhibit growth of CCL64 cells. Epitope-tagged TGF- β 1 activity was present at dilutions of over 1:250, and

could be neutralized with the anti-TGF- β blocking antibody, 2G7, thereby demonstrating that the inhibition was specific to the TGF- β protein or fusion protein.

FIG 6A illustrates the design of TGF- β 1 with an N-terminal FLAG tag (NFLAG-TGF- β 1). The FLAG tag (boxed sequence) was inserted immediately following the cleavage site (indicated by arrow). Amino acid sequence shown above
5 nucleotide sequence in single letter code.

FIG 6B shows the expression levels of secreted TGF- β 1 from transfected Cos1 cells (left panel). Blot probed with a rabbit polyclonal anti-human TGF- β 1 antibody (Promega). For the NFLAG-TGF- β 1 transfection, the ratio of cleaved,
10 mature peptide (m) to unprocessed pro-peptide (p) is less than that for the untagged ligand (WT-TGF- β 1). In the middle panel, a Western blot illustrates that anti-FLAG monoclonal antibody (clone M2) specifically detects the FLAG-tagged TGF- β 1 but not the untagged ligand. Finally, the right panel of FIG 6B illustrates that N-FLAG-TGF- β 1 shows smad2 phosphorylation activity, based on the signal labeled "anti-P-
15 smad2. rh-TGF- β 1, recombinant human TGF- β 1 at 2.5 ng/mL; levels of total smad 2 protein are unchanged.

FIG 6C shows portions of the sequences of N+5FLAG-TGF- β 1 and N+5HA-TGF- β 1. Only the region in the vicinity of the cleavage site is shown; the numbering is different than in the attached sequence listing and refers to the position
20 of the illustrated residues in the plasmid. The epitope tag sequences are boxed. Arrows indicate cleavage site that separates LAP from the mature TGF- β peptide. Note the localized concentration of basic amino acids in the region immediately upstream of the cleavage site (RHRR).

FIG 6D shows the levels of secreted TGF- β 1 detected by immunoblot
25 analysis using a polyclonal anti-human TGF- β 1 antibody (first four lanes) and the epitope tags (probed with anti-FLAG antibodies in the middle pair of lanes; probed with anti-HA antibodies right pair of lanes) for each of the indicated constructs. The ratio of mature to unprocessed ligand is essentially equivalent for each construct. Antibodies against each of the epitope tags detect only the respective tagged ligand
30 and not the un-tagged molecule.

FIG 7 illustrates activity of TGF- β fusion proteins. Mv1Lu cells were treated with conditioned medium from transfected cos-1 cells and inhibition of growth was assessed by incorporation of [3 H] thymidine. Results are shown in FIG 7A. Solid bars, N+5FLAG-TGF- β 1; first set of grey bars in each half of the graph, N+5HA-TGF- β 1; open bars, WT-TGF- β 1; second set of grey bars in each half of the graph, Empty vector. - and + indicate, respectively, absence or presence of TGF- β neutralizing antibody.

FIG 7B is a pair of western blots illustrating levels of phospho-smad2 (upper panel) and smad2 (lower panel). Mv1Lu cells were treated with conditioned media (diluted 1:20) from Cos1 cells transfected with plasmids encoding untagged TGF- β 1 (WT-TGF- β 1), N+5FLAG-TGF- β 1, N+5HA-TGF- β 1 or empty vector. Rh-TGF- β 1, recombinant human TGF- β 1 (2.5 ng/mL). - and + indicate, respectively, the absence or presence of TGF- β neutralizing antibody. To generate the signals on the lower panel, the blot was stripped and re-probed using an antibody directed against smad 2.

FIG 7C illustrates the quantitative results of a sandwich ELISA to detect tagged TGF- β 1 ligands in conditioned media. Solid bars, FLAG-tagged TGF- β 1; stippled bars, HA-tagged TGF- β 1; gray bars, untagged (WT) TGF- β 1 in a FLAG-TGF- β 1 SELISA; open bars, untagged (WT) TGF- β 1 in a HA-TGF- β 1 SELISA. "No 1st," negative control where primary antibody (anti-FLAG or anti-HA) was omitted. Dilutions refer to dilutions of conditioned media.

SEQUENCE LISTING

The nucleic and amino acid sequences listed in the accompanying sequence listing are shown using standard letter abbreviations for nucleotide bases, and three letter code for amino acids, as defined in 37 C.F.R. 1.822. Only one strand of each nucleic acid sequence is shown, but the complementary strand is understood as included by any reference to the displayed strand. In the accompanying sequence listing:

SEQ ID NOs: 1 and 2 show the nucleic acid sequences of the flanking primers used to introduce the FLAG epitope tag into TGF- β 1 using 2-step PCR mutagenesis technique.

5 SEQ ID NOs: 3 and 4 show the nucleic acid sequences of the primers used to construct the N-terminally tagged-TGF- β 1 fusion (FLAG- β 1).

SEQ ID NOs: 5 and 6 show the nucleic acid sequences of the primers used to insert the FLAG peptide tag between amino acids 11 and 12 of the mature TGF- β 1.

10 SEQ ID NO: 7 shows the amino acid sequence of the FLAG peptide tag.
SEQ ID NO: 8 shows the nucleotide and amino acid sequences of the precursor of the NFLAG-TGF- β 1 fusion.

15 SEQ ID NO: 9 shows the amino acid sequence of the precursor of the N FLAG- β 1 fusion. The cleavage site for maturation of the fusion is between amino acid residues 278 and 279, resulting in a mature NFLAG-TGF- β 1 fusion of 120 amino acid residues.

SEQ ID NO: 10 shows the nucleic acid sequence of the mature NFLAG- β 1 fusion.

20 SEQ ID NO: 11 shows the amino acid sequence of the mature NFLAG- β 1 fusion.

SEQ ID NO: 12 shows the nucleotide and amino acid sequences of the precursor of the fusion with the FLAG peptide tag between amino acids 11 and 12 of the mature TGF- β 1.

25 SEQ ID NO: 13 shows the amino acid sequence of the precursor of the fusion with the FLAG peptide tag between amino acids 11 and 12 of the mature TGF- β 1. The cleavage site for maturation of the fusion is between amino acid residues 278 and 279, resulting in a mature 11/12FLAG- β 1 fusion of 120 amino acid residues.

SEQ ID NO: 14 shows the nucleic acid sequence of the mature 11/12FLAG- β 1 fusion.

SEQ ID NO: 15 shows the amino acid sequence of the mature 11/12FLAG- β 1 fusion.

SEQ ID NO: 16 shows the nucleic acid and deduced amino acid sequence of the precursor murine N+5FLAG TGF- β 1 (N+5FLAG- β 1) fusion.

5 SEQ ID NO: 17 shows the amino acid sequence of the precursor murine N+5FLAG TGF- β 1 (N+5FLAG- β 1) fusion.

SEQ ID NOs: 18 and 19 show forward and reverse primers (respectively) used in construction of the N+5FLAG TGF- β 1 (N+5FLAG- β 1) fusion.

10 SEQ ID NO: 20 shows the nucleic acid and deduced amino acid sequence of the precursor N+5HA-TGF- β 1 (N+5HA- β 1) construct.

SEQ ID NO: 21 shows the amino acid sequence of the precursor N+5HA-TGF- β 1 (N+5HA- β 1) fusion.

SEQ ID NOs: 22 and 23 show forward and reverse primers (respectively) used in construction of the N+5HA-TGF- β 1 (N+5HA- β 1) fusion.

15 SEQ ID NO: 24 shows the nucleic acid and deduced protein sequence of the precursor murine N+5FLAG TGF- β 2 (MN5FLAGb2) fusion.

SEQ ID NO: 25 shows the amino acid sequence of the precursor N+5FLAG-TGF- β 2 (MN5FLAGb2) fusion.

20 SEQ ID NO: 26 shows the nucleic acid and deduced protein sequence of the precursor murine N+5HA TGF- β 2 (MN5HAb2) fusion.

SEQ ID NO: 27 shows the amino acid sequence of the precursor N+5HA-TGF- β 2 (MN5HAb2) fusion.

SEQ ID NO: 28 shows the nucleic acid and deduced protein sequence of the precursor murine N+5FLAG TGF- β 3 (MN5FLAGb3) fusion.

25 SEQ ID NO: 29 shows the amino acid sequence of the precursor N+5FLAG-TGF- β 3 (MN5FLAGb3) fusion.

SEQ ID NO: 30 shows the nucleic acid and deduced protein sequence of the precursor murine N+5HA TGF- β 3 (MN5HAb3) fusion.

30 SEQ ID NO: 31 shows the amino acid sequence of the precursor N+5HA-TGF- β 3 (MN5HAb3) fusion.

SEQ ID NO: 32 shows the nucleic acid and deduced protein sequence of the precursor porcine active N+5FLAG TGF- β 1 (actN5FLAGb1) fusion.

SEQ ID NO: 33 shows the amino acid sequence of the precursor porcine active N+5FLAG-TGF- β 1 (actN5FLAGb1) fusion.

5 SEQ ID NO: 34 shows the nucleic acid and deduced protein sequence of the precursor porcine latent N+5FLAG TGF- β 1 (latN5FLAGb1) fusion.

SEQ ID NO: 35 shows the amino acid sequence of the precursor porcine latent N+5FLAG-TGF- β 1 (latN5FLAGb1) fusion.

Sub B1 10 ~~SEQ ID NO: 36 shows the nucleic acid and deduced protein sequence of the precursor porcine latent N+5FLAG TGF- β 1 (actN5HAb1) fusion.~~

SEQ ID NO: 37 shows the amino acid sequence of the precursor N+5HA-TGF- β 1 (actN5HAb1) fusion.

Sub B2 ~~SEQ ID NO: 38 shows the nucleic acid and deduced protein sequence of the precursor porcine active N+5FLAG TGF- β 1 (latN5HAb1) fusion.~~

15 SEQ ID NO: 39 shows the amino acid sequence of the precursor N+5HA-TGF- β 1 (latN5HAb1) fusion.

DETAILED DESCRIPTION OF THE DISCLOSURE

I. Abbreviations

- 20 LAP: latency associated peptide
NMR: nuclear magnetic resonance
DTT: dithiothreitol
ELISA: enzyme linked immunoassay
SELISA: sandwich ELISA
25 TGF: transforming growth factor

II. Terms

Unless otherwise noted, technical terms are used according to conventional usage. Definitions of common terms in molecular biology may be found in

- 30 Benjamin Lewin, *Genes V*, published by Oxford University Press, 1994 (ISBN 0-19-854287-9); Kendrew *et al.* (eds.), *The Encyclopedia of Molecular Biology*, published

by Blackwell Science Ltd., 1994 (ISBN 0-632-02182-9); A.D. Smith *et al.* (eds.), *Oxford Dictionary of Biochemistry and Molecular Biology*, published by Oxford Univ. Press, 1997 (ISBN 0-19-854768-4); and Robert A. Meyers (ed.), *Molecular Biology and Biotechnology: a Comprehensive Desk Reference*, published by VCH Publishers, Inc., 1995 (ISBN 1-56081-569-8).

In order to facilitate review of the various embodiments, the following explanations of terms are provided:

Associating: A process of reversibly bringing two chemical entities (such as biological macromolecules, *e.g.*, proteins) together to form a complex, usually where the association involves specific interactions between the macromolecules. Two biological macromolecules may associate with each other automatically, for instance during synthesis of the molecules, or may require one or more chaperon-like molecules to mediate the association process.

Cleaving: To split a chemical bond, such as a chemical linkage between the subunits of a biopolymer. Thus, breaking a peptide bond in a oligo- or polypeptide (*e.g.*, in a protein) is a cleavage event. Cleaving may be relatively non-specific, for instance though a purely chemical process of acidic hydrolysis. In some instances, cleavages are mediated by a biochemical or biological process, for instance through the process of an enzyme such as a protease (in the case of a peptide cleavage) or a nuclease (in the case of cleavage of a poly-nucleotide). As will be recognized by those of ordinary skill in the art, enzyme-based cleavage is often a specific event, and for instance may be influenced by the primary and/or secondary structure of the molecule (*e.g.*, biopolymer) that is being cleaved.

A specific class of cleavage events are those involved in maturation of precursor proteins, for instance the proteolytic cleavage ("removal") of a pro-region portion of a precursor (*e.g.*, a latent) protein.

Epitope tags are short stretches of amino acids to which a specific antibody can be raised, which in some embodiments allows one to specifically identify and track the tagged protein that has been added to a living organism or to cultured cells. Detection of the tagged molecule can be achieved using a number of different

techniques. Examples of such techniques include: immunohistochemistry, immunoprecipitation, flow cytometry, immunofluorescence microscopy, ELISA, immunoblotting ("western"), and affinity chromatography. Examples of useful epitope tags include FLAG, T7, HA (hemagglutinin) and myc. The FLAG tag
5 (DYKDDDDK) was used in some particular examples disclosed herein because high quality reagents are available to be used for its detection. Epitope tags can of course have one or more additional functions, beyond recognition by an antibody.

Functionalized: To add a function to a molecule, such as a protein, that is not a native function of that molecule. A **functionalizing peptide portion** is a
10 portion of a protein (*e.g.*, a peptide or protein domain), for instance a part of a fusion protein, that provides a non-native function to the fusion. Functions that can be added in this manner include (but are not limited to) identification (particularly distinction from the native subject protein), targeting, translocation, and distinct biological function (such as addition of an active protein domain).

Fusion protein: A protein comprising two amino acid sequences that are not found joined together in nature. **Functionalized TGF- β fusion proteins**
15 specifically comprise at least (1) a sequence of a mature region of a TGF- β family protein and (2) a functionalizing peptide portion placed at either end of or within the coding region of the mature TGF- β family protein. Such functionalized TGF- β
20 fusion proteins can additionally include other protein elements, such as one or more additional functionalizing peptide portion, a pro-region of a TGF- β family protein (which may assist in folding, assembly, and/or secretion of the fusion protein), and/or a linker between any of such protein portions ("domains").

Injectable composition: A pharmaceutically acceptable fluid composition
25 comprising at least one active ingredient, *e.g.*, a functionalized TGF- β family protein fusion. The active ingredient is usually dissolved or suspended in a physiologically acceptable carrier, and the composition can additionally comprise minor amounts of one or more non-toxic auxiliary substances, such as emulsifying agents, preservatives, and pH buffering agents and the like. Such injectable compositions
30 that are useful for use with the fusion proteins of this disclosure are conventional; appropriate formulations are well known to those of ordinary skill in the art.

- Isolated:** An "isolated" biological component (such as a nucleic acid molecule, protein or organelle) has been substantially separated or purified away from other biological components in the cell of the organism in which the component naturally occurs, *i.e.*, other chromosomal and extra-chromosomal DNA and RNA, proteins and organelles. Nucleic acids and proteins that have been "isolated" include nucleic acids and proteins purified by standard purification methods. The term also embraces nucleic acids and proteins prepared by recombinant expression in a host cell, as well as chemically synthesized biopolymers.
- Linker:** A peptide, usually between two and about 150 amino acid residues in length, that serves to join two protein domains in a multi-domain (multi-part) fusion protein. Examples of specific linkers can be found, for instance, in Hennecke *et al.* (*Protein Eng.* 11:405-410, 1998); and U.S. Patent Nos. 5,767,260 and 5,856,456.
- Linkers may be repetitive or non-repetitive. One classical repetitive linker used in the production of fusion proteins is the (Gly₄Ser)₃ (or (GGGS)₃ or (G₄S)₃) linker. More recently, non-repetitive linkers have been produced, and methods for the random generation of such linkers are known (Hennecke *et al.*, *Protein Eng.* 11:405-410, 1998). In addition, linkers may be chosen to have more or less secondary character (*e.g.*, helical character, U.S. Patent No. 5,637,481) depending on the conformation desired in the final fusion protein. The more secondary character a linker possesses, the more constrained the structure of the final fusion protein will be. Therefore, substantially flexible linkers that are substantially lacking in secondary structure allow flexion of the fusion protein at the linker.
- Nucleotide:** "Nucleotide" includes, but is not limited to, a monomer that includes a base linked to a sugar, such as a pyrimidine, purine or synthetic analogs thereof, or a base linked to an amino acid, as in a peptide nucleic acid (PNA). A nucleotide is one monomer in a polynucleotide. A nucleotide sequence refers to the sequence of bases in a polynucleotide.
- Oligonucleotide:** An oligonucleotide is a plurality of joined nucleotides joined by native phosphodiester bonds, usually between about 6 and about 300

nucleotides in length. The term "oligonucleotide analog" refers to moieties that function similarly to oligonucleotide but have non-naturally occurring portions. For example, oligonucleotide analogs can contain altered sugar moieties or inter-sugar linkages, such as a phosphorothioate oligodeoxynucleotide. Functional analogs of naturally occurring polynucleotides can bind to RNA or DNA, and include peptide nucleic acid (PNA) molecules.

Particular oligonucleotides and oligonucleotide analogs can include linear sequences up to about 300 nucleotides in length, for example a sequence (such as DNA or RNA) that is at least 6 bases, for example at least 8, 10, 15, 20, 25, 30, 35, 40, 45, 50, 100, 200, or even 300 bases long, or from about 6 to about 100 bases, for example about 10-50 bases, such as 12, 15, 20, 30, or 40 bases.

Open reading frame: A series of nucleotide triplets (codons) coding for amino acids without any internal termination codons. These sequences are usually translatable into a peptide.

Operably linked nucleic acid: A first nucleic acid sequence is operably linked with a second nucleic acid sequence when the first nucleic acid sequence is placed in a functional relationship with the second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Generally, operably linked DNA sequences are contiguous and, where necessary to join two protein-coding regions, in the same reading frame.

ORF (open reading frame): A series of nucleotide triplets (codons) coding for amino acids without any internal termination codons. These sequences are usually translatable into a peptide.

Parenteral: Administered outside of the intestine, *e.g.*, not via the alimentary tract. Generally, parenteral formulations are those that will be administered through any possible mode except ingestion. This term especially refers to injections, whether administered intravenously, intrathecally, intramuscularly, intraperitoneally, or subcutaneously, and various surface applications including intranasal, intradermal, and topical application, for instance.

Peptide: Any compound containing two or more amino-acid residues joined by amide bonds, formed from the carboxyl group of one residue and the amino group of the next. The broad term "peptide" includes oligopeptides, polypeptides, and proteins.

- 5 **Peptide Nucleic Acid (PNA):** An oligonucleotide analog with a backbone comprised of monomers coupled by amide (peptide) bonds, such as amino acid monomers joined by peptide bonds.

10 **Peptide tag:** A peptide sequence that is attached (for instance through genetic engineering) to another peptide or a protein, to provide a function to the resultant fusion. Peptide tags are usually relatively short in comparison to a protein to which they are fused; by way of example, peptide tags are four or more amino acids in length, such as 5, 6, 7, 8, 9, 10, 15, 20, or 25 or more amino acids. Usually a peptide tag will be no more than about 100 amino acids in length, and may be no more than about 75, no more than about 50, no more than about 40, or no more than
15 about 30.

 In preferred embodiments, the addition of the functionalizing peptide tag will not substantially reduce a native function of the TGF- β molecule to which it is attached. Thus, in particular examples, the addition of a functionalizing peptide tag will reduce a function of the TGF- β molecule by no more than about 50%, about
20 40%, about 30%, about 20%, or about 10%. In certain specific examples, the activity of the so-modified TGF- β molecule will be reduced by no more than about 5%, about 3%, about 1%, or may in fact not be reduced at all. Native TGF- β activity can be measured using any of various known

 Peptide tags confer one or more different functions to a fusion protein
25 (thereby "functionalizing" that protein), and such functions can include (but are not limited to) antibody binding (an epitope tag), purification, translocation, targeting, and differentiation (*e.g.*, from a native protein). In addition, a recognition site for a protease, for which a binding antibody is known, can be used as a specifically cleavable epitope tag. The use of such a cleavable tag can provide selective
30 cleavage and activation of a protein (*e.g.*, by replacing the cleavage site in TGF- β 1 with that for pro-caspase 3. Alternatively the system developed by in the Dowdy

laboratory (Vecero-Akbani *et al.*, *Nat Med.* 5:29-33, 1999) could be use to provide specificity of such cleavage and activation.

Detection of the tagged molecule can be achieved using a number of different techniques. These include: immunohistochemistry, immunoprecipitation,
5 flow cytometry, immunofluorescence microscopy, ELISA, immunoblotting ("western"), and affinity chromatography.

Epitope tags add a known epitope (antibody binding site) on the subject protein, to provide binding of a known and often high-affinity antibody, and thereby allowing one to specifically identify and track the tagged protein that has been added
10 to a living organism or to cultured cells. Examples of epitope tags include the myc, T7, GST, GFP, HA (hemagglutinin) and FLAG tags. The first four examples are epitopes derived from existing molecules. In contrast, FLAG is a synthetic epitope tag designed for high antigenicity (see, *e.g.*, U.S. Patent Nos. 4,703,004 and 4,851,341).

Purification tags are used to permit easy purification of the tagged protein, such as by affinity chromatography. A well-known purification tag is the hexa-histidine (6x His) tag, literally a sequence of six histidine residues. The 6x His protein purification system is available commercially from QIAGEN (Valencia, CA), under the name of QIAexpress®.

One representative translocation/transduction peptide (which serves to facilitate translocation of a protein into a cell) is the Tat₄₉₋₅₇ (RKKRRQRRR) fragment from the human immunodeficiency virus (HIV). See, for instance, Vecero-Akbani *et al.* (*Methods Enzymol.* 322:508-521, 2000); Falnes *et al.* (*Biochem.* 40:4349-4358, 2001); and Becker-Hapak *et al.* (*Methods* 24:247-256,
25 2001). Antibodies are available and can be produced that recognize this peptide, and so it can be viewed as a multi-functional peptide.

A single tag peptide can serve more than one purpose; any attached tag, for instance, will increase the molecular weight of the fusion protein and thereby permit differentiation between the tagged and native proteins. Antibodies specific for an
30 "epitope tag" can be used to construct an immunoaffinity column, thus permitting an epitope tag to be used for purification of the tagged protein. Likewise, in some

instances monoclonal antibodies specific for a purification tag are available (*e.g.*, anti-6x His peptide monoclonal antibodies, which are available through QIAGEN or CLONTECH, Palo Alto, CA).

Pharmaceutically acceptable carriers: The pharmaceutically acceptable carriers useful in the disclosed compositions and methods are conventional. *Remington's Pharmaceutical Sciences*, by E. W. Martin, Mack Publishing Co., Easton, PA, 15th Edition (1975), describes compositions and formulations suitable for pharmaceutical delivery of the fusion proteins herein disclosed.

In general, the nature of the carrier will depend on the particular mode of administration being employed. For instance, parenteral formulations usually comprise injectable fluids that include pharmaceutically and physiologically acceptable fluids such as water, physiological saline, balanced salt solutions, aqueous dextrose, glycerol or the like as a vehicle. For solid compositions (*e.g.*, powder, pill, tablet, or capsule forms), conventional non-toxic solid carriers can include, for example, pharmaceutical grades of mannitol, lactose, starch, or magnesium stearate. In addition to biologically-neutral carriers, pharmaceutical compositions to be administered can contain minor amounts of non-toxic auxiliary substances, such as wetting or emulsifying agents, preservatives, and pH buffering agents and the like, for example sodium acetate or sorbitan monolaurate.

Pharmacodynamics is the branch of pharmacology dealing with the effects of drugs on the body, *i.e.*, with the physiological, therapeutic, and toxicological responses to drugs with particular regard to the extent, mechanism, and time course of such effects on living organisms (see, Goodman and Gilman, "*The Pharmacological Basis of Therapeutics*," Macmillan Publ. Co., 1985, LC # 85-15356, Chapter 2, pages 34-65). The study of pharmacodynamics may include measurement or determination of the cellular site of drug action, dose-response relationship(s), structure-activity relationship(s), or quantitation of drug-receptor interactions, for example.

Pharmacokinetics is the branch of pharmacology dealing quantitatively with the movement of drugs (*e.g.*, proteins) within the body, *i.e.*, with the absorption (uptake into the system), bioavailability (fraction of the drug absorbed as such into

systemic circulation), distribution (movement around the available volume of the body), metabolism (conversion of the drug into non-active form(s)), and elimination/clearance (the ability of a body to eliminate the drug from the system) of drugs. Each of these factors influence the magnitude of a drug effect(s), and how much of the drug found at sites at different times after its application to a system (see, Levine, "*Pharmacology: Drug Actions and Reactions*," Little, Brown and Co., Ltd., 1983, ISBN 0-316-5222-8, Chapter 10, pages 211-247, and Goodman and Gilman, "*The Pharmacological Basis of Therapeutics*," Macmillan Publ. Co., 1985, LC # 85-15356, Chapter 1, pages 3-34).

10 **Pharmacological properties of a drug:** The properties of a drug (any non-food chemical agent, such as a protein, that affects a living organism) in a subject, in particular as relates to the use of drugs to treat, cure, prevent, diagnose, or prognose disease. Pharmacological properties include the pharmacodynamic and pharmacokinetic properties, for instance.

15 **Polymorphism:** Variant in a sequence of a gene. Polymorphisms can be those variations (nucleotide sequence differences) that, while having a different nucleotide sequence, produce functionally equivalent gene products, such as those variations generally found between individuals, different ethnic groups, geographic locations. The term polymorphism also encompasses variations that produce gene products with altered function, *i.e.*, variants in the gene sequence that lead to gene products that are not functionally equivalent. This term also encompasses variations that produce no gene product, an inactive gene product, or increased gene product. The term polymorphism may be used interchangeably with allele or mutation, unless context clearly dictates otherwise.

25 Polymorphisms can be referred to, for instance, by the nucleotide position at which the variation exists, by the change in amino acid sequence caused by the nucleotide variation, or by a change in some other characteristic of the nucleic acid molecule that is linked to the variation (*e.g.*, an alteration of a secondary structure such as a stem-loop, or an alteration of the binding affinity of the nucleic acid for associated molecules, such as polymerases, RNases, and so forth).

30

Probes and primers: A probe comprises an isolated nucleic acid attached to a detectable label or other reporter molecule. Typical labels include radioactive isotopes, enzyme substrates, co-factors, ligands, chemiluminescent or fluorescent agents, haptens, and enzymes. Methods for labeling and guidance in the choice of labels appropriate for various purposes are discussed, *e.g.*, in Sambrook *et al.* (In *Molecular Cloning: A Laboratory Manual*, CSHL, New York, 1989) and Ausubel *et al.* (In *Current Protocols in Molecular Biology*, John Wiley & Sons, New York, 1998).

Primers are short nucleic acid molecules, for instance DNA oligonucleotides about 6 nucleotides or more in length. Longer DNA oligonucleotides may be about 10, 15, 20, 25, 30 or 50 nucleotides or more in length. Primers can be annealed to a complementary target DNA strand by nucleic acid hybridization to form a hybrid between the primer and the target DNA strand, and then the primer extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification of a nucleic acid sequence, *e.g.*, by the polymerase chain reaction (PCR) or other nucleic-acid amplification methods known in the art. Other examples of amplification include, but are not limited to, strand displacement amplification, as disclosed in U.S. Patent No. 5,744,311; transcription-free isothermal amplification, as disclosed in U.S. Patent No. 6,033,881; repair chain reaction amplification, as disclosed in WO 90/01069; ligase chain reaction amplification, as disclosed in EP-A-320 308; gap filling ligase chain reaction amplification, as disclosed in 5,427,930; and NASBATM RNA transcription-free amplification, as disclosed in U.S. Patent No. 6,025,134.

Methods for preparing and using nucleic acid probes and primers are described, for example, in Sambrook *et al.* (In *Molecular Cloning: A Laboratory Manual*, CSHL, New York, 1989), Ausubel *et al.* (ed.) (In *Current Protocols in Molecular Biology*, John Wiley & Sons, New York, 1998), and Innis *et al.* (*PCR Protocols, A Guide to Methods and Applications*, Academic Press, Inc., San Diego, CA, 1990). Amplification primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, © 1991, Whitehead Institute for Biomedical Research, Cambridge,

MA). One of ordinary skill in the art will appreciate that the specificity of a particular probe or primer increases with its length. Thus, for example, a primer comprising 30 consecutive nucleotides of a TGF- β -encoding nucleotide or flanking region thereof (such as a "TGF- β 1 primer" or "TGF- β 1 probe") will anneal to a target sequence with a higher specificity than a corresponding primer of only 15 nucleotides. Thus, in order to obtain greater specificity, probes and primers can be selected that comprise at least 20, 25, 30, 35, 40, 45, 50 or more consecutive nucleotides of the target nucleotide sequences.

Protein: A biological molecule expressed by an encoding nucleic acid molecule (*e.g.*, a gene) and comprised of amino acids. Proteins are a subset of the broader molecular class "peptide."

Purified: The term purified does not require absolute purity; rather, it is intended as a relative term. Thus, for example, a "purified" fusion protein preparation is one in which the fusion protein is more enriched than the protein is in its generative environment, for instance within a cell or in a biochemical reaction chamber. Preferably, a preparation of fusion protein is purified such that the fusion protein represents at least 50% of the total protein content of the preparation.

Recombinant nucleotide: A recombinant nucleic acid molecule is one that has a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two otherwise separated segments of sequence. This artificial combination can be accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, *e.g.*, by genetic engineering techniques.

Similarly, a recombinant protein is one encoded for by a recombinant nucleic acid molecule.

Sequence identity: The similarity between two nucleic acid sequences, or two amino acid sequences, is expressed in terms of the similarity between the sequences, otherwise referred to as sequence identity. Sequence identity is frequently measured in terms of percentage identity (or similarity or homology); the higher the percentage, the more similar the two sequences are.

Methods of alignment of sequences for comparison are well known in the art. Various programs and alignment algorithms are described in: Smith and Waterman (*Adv. Appl. Math.* 2: 482, 1981); Needleman and Wunsch (*J. Mol. Biol.* 48: 443, 1970); Pearson and Lipman (*PNAS. USA* 85: 2444, 1988); Higgins and Sharp
5 (*Gene*, 73: 237-244, 1988); Higgins and Sharp (*CABIOS* 5: 151-153, 1989); Corpet
et al. (*Nuc. Acids Res.* 16: 10881-10890, 1988); Huang *et al.* (*Comp. Appls Biosci.*
8: 155-165, 1992); and Pearson *et al.* (*Meth. Mol. Biol.* 24: 307-31, 1994). Altschul
et al. (*Nature Genet.*, 6: 119-129, 1994) presents a detailed consideration of
sequence alignment methods and homology calculations.

10 The alignment tools ALIGN (Myers and Miller, *CABIOS* 4:11-17, 1989) or
LFASTA (Pearson and Lipman, 1988) may be used to perform sequence
comparisons (Internet Program © 1996, W. R. Pearson and the University of
Virginia, "fasta20u63" version 2.0u63, release date December 1996). ALIGN
compares entire sequences against one another, while LFASTA compares regions of
15 local similarity. These alignment tools and their respective tutorials are available on
the Internet at the NCSA Website. Alternatively, for comparisons of amino acid
sequences of greater than about 30 amino acids, the "Blast 2 sequences" function
can be employed using the default BLOSUM62 matrix set to default parameters,
(gap existence cost of 11, and a per residue gap cost of 1). When aligning short
20 peptides (fewer than around 30 amino acids), the alignment should be performed
using the "Blast 2 sequences" function, employing the PAM30 matrix set to default
parameters (open gap 9, extension gap 1 penalties). The BLAST sequence
comparison system is available, for instance, from the NCBI web site; see also
Altschul *et al.*, *J. Mol. Biol.* 215:403-410, 1990; Gish. & States, *Nature Genet.*
25 3:266-272, 1993; Madden *et al.* *Meth. Enzymol.* 266:131-141, 1996; Altschul *et al.*,
Nucleic Acids Res. 25:3389-3402, 1997; and Zhang & Madden, *Genome Res.* 7:649-
656, 1997.

Orthologs of proteins are typically characterized by possession of greater
than 75% sequence identity counted over the full-length alignment with the amino
30 acid sequence of specific protein using ALIGN set to default parameters. Proteins
with even greater similarity to a reference sequence will show increasing percentage

identities when assessed by this method, such as at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, or at least 98% sequence identity. In addition, sequence identity can be compared over the full length of one or both binding domains of the disclosed fusion proteins.

5 When significantly less than the entire sequence is being compared for sequence identity, homologous sequences will typically possess at least 80% sequence identity over short windows of 10-20 amino acids, and may possess sequence identities of at least 85%, at least 90%, at least 95%, or at least 99% depending on their similarity to the reference sequence. Sequence identity over such
10 short windows can be determined using LFASTA; methods are described at the NCSA Website. One of skill in the art will appreciate that these sequence identity ranges are provided for guidance only; it is entirely possible that strongly significant homologs could be obtained that fall outside of the ranges provided.

 An alternative indication that two nucleic acid molecules are closely related
15 is that the two molecules hybridize to each other under stringent conditions. Stringent conditions are sequence-dependent and are different under different environmental parameters. Generally, stringent conditions are selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic
20 strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Conditions for nucleic acid hybridization and calculation of stringencies can be found in Sambrook *et al.* (In *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor, New York, 1989) and Tijssen (*Laboratory Techniques in Biochemistry and Molecular Biology Part I*, Ch. 2, Elsevier, New York, 1993).
25 Nucleic acid molecules that hybridize under stringent conditions to the disclosed bispecific fusion protein sequences will typically hybridize to a probe based on either the entire fusion protein encoding sequence, an entire binding domain, or other selected portions of the encoding sequence under wash conditions of 0.2 x SSC, 0.1% SDS at 65°C.

30 Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences, due to the degeneracy of the

genetic code. It is understood that changes in nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that each encode substantially the same protein.

Specific binding agent: An agent that binds substantially only to a defined target. Thus a peptide-specific binding agent binds substantially only the defined peptide, or a peptide region within a protein, such as a fusion protein. As used herein, the term "[X] specific binding agent," where [X] refers to a specific protein or peptide, includes anti-[X] antibodies (and functional fragments thereof) and other agents (such as soluble receptors) that bind substantially only to [X].

Antibodies may be produced using standard procedures described in a number of texts, including Harlow and Lane (*Antibodies, A Laboratory Manual*, CSHL, New York, 1988). The determination that a particular agent binds substantially only to the target protein or peptide may readily be made by using or adapting routine procedures. One suitable *in vitro* assay makes use of the Western blotting procedure (described in many standard texts, including Harlow and Lane, *Antibodies, A Laboratory Manual*, CSHL, New York, 1988). Western blotting may be used to determine that a given protein binding agent, such as an anti-TGF- β family protein monoclonal antibody, binds substantially only to the specified TGF- β family protein.

Shorter fragments of antibodies can also serve as specific binding agents. For instance, FAbs, Fvs, and single-chain Fvs (SCFvs) that bind to a protein or peptide [X] would be [X]-specific binding agents. These antibody fragments are defined as follows: (1) FAb, the fragment which contains a monovalent antigen-binding fragment of an antibody molecule produced by digestion of whole antibody with the enzyme papain to yield an intact light chain and a portion of one heavy chain; (2) FAb', the fragment of an antibody molecule obtained by treating whole antibody with pepsin, followed by reduction, to yield an intact light chain and a portion of the heavy chain; two FAb' fragments are obtained per antibody molecule; (3) (FAb')₂, the fragment of the antibody obtained by treating whole antibody with the enzyme pepsin without subsequent reduction; (4) F(Ab')₂, a dimer of two FAb' fragments held together by two disulfide bonds; (5) Fv, a genetically engineered

fragment containing the variable region of the light chain and the variable region of the heavy chain expressed as two chains; and (6) single chain antibody ("SCA"), a genetically engineered molecule containing the variable region of the light chain, the variable region of the heavy chain, linked by a suitable polypeptide linker as a
5 genetically fused single chain molecule. Methods of making these fragments are routine.

Subject: Living multi-cellular vertebrate organisms, a category that includes both human and non-human mammals.

TGF- β family of proteins: A family of secreted signaling molecules
10 involved in a number of cellular and developmental processes in eukaryotic cells, including inflammation, immune surveillance, and neoplasia. Members of the TGF- β family of proteins include, but are not limited to: TGF- β 2, TGF- β 3, TGF- β 1, TGF- β 4 (chicken), TGF- β 5 (*Xenopus*), GDF-9 (mouse/human), BMP-16/nodal (mouse), Fugacin (*Xenopus*), BMP3, Sumitomo-BIP/GDF-10 (mouse), ADMP
15 (*Xenopus*), BMP-9, Dorsalin-1 (Chicken), BMP-10, BMP-13/GDF-6 (mouse), Radar (Zebrafish), GDF-1/CDMP-1 (mouse/human), BMP-12/GDF-7 (mouse), BMP-5, BMP-6, BMP-7/OP-1, BMP-8/OP-2, PC8/OP-3 (mouse), 60A (*Drosophila*), BMP-2, BMP-4, Decapentaplegic (*Drosophila*), Vg-1 (*Xenopus*), Univin (sea urchin), Vgr-2/GDF-3, GDF-1, Screw (*Drosophila*), BMP-11, GDF-8, Activin β C,
20 Activin β D (*Xenopus*), Activin β E, BMP-14/GDF-12, Activin β A, Activin β B, GDF-14, Mullerian inhibiting substance, and α -inhibin. The phylogenetic relationship between these proteins is shown in FIG 1.

The term "TGF- β family protein function" includes all functions that are associated with a TGF- β family protein, including for instance secondary folding of
25 each TGF- β monomer, tertiary association between the members of the multimeric (e.g., homodimeric) TGF- β complex, maturation by cleavage and/or removal of the pro-region (LAP), secretion of the protein from the cell in which it was translated, specific receptor binding, and down-stream activities that result from the binding of a TGF- β family ligand protein with its cognate receptor(s). Such downstream
30 activities include (depending on the TGF- β family member examined and the system

used), for instance, regulation of cell growth (proliferation), stimulation of cell growth or proliferation, stimulation of cell differentiation, inhibition of cell growth or proliferation, regulation of cytokine production, induction of cellular differentiation, cell cycle inhibition, control of adhesion molecule expression, stimulation of angiogenesis, induction of leukocyte chemotaxis, induction of apoptosis, suppression of lymphocyte activation, suppression of inflammation, enhances wound healing by mechanisms including , stimulation of synthesis of matrix proteins, regulation of immunoglobulin production, including isotype switch recombination, and suppression of tumorigenesis.

10 Fusions of the current disclosure maintain substantial TGF- β biological activity, by which it is meant that the fusion protein maintains at least one biological activity at a level of at least approximately 50% of the native equivalent TGF- β protein. In specific embodiments, the fusion protein will maintain a greater level of one or more specific TGF- β biological activities, such as at least 60%, at least 70%,
15 at least 80%, or at least 90% or more of the native activity. In certain embodiments, a functionalized TGF- β fusion protein of the disclosure will display a specific TGF- β biological activity equal to or greater than that observed with the equivalent native TGF- β protein, for instance at least 100% of the native activity, or 105%, 110%, 120%, or even 150% of native (non-fused) TGF- β .

20 Different members of the TGF- β family have different biological specificities and activities. Specificities of the listed TGF- β family proteins are known to one of ordinary skill in the art. See, for instance, Doetschman, *Lab. Anim. Sci.* 49:137-143, 1999; Letterio and Roberts, *Annu. Rev. Immunol.* 16:137-61:137-161, 1998; Wahl, *J. Exp. Med.* 180:1587-1590, 1994; Letterio and
25 Roberts, *J. Leukoc. Biol.* 59:769-774, 1996; Piek *et al.*, *FASEB J.* 13:2105-2124, 1999; Heldin *et al.*, *Nature* 390:465-471, 1979; and De Caestecker *et al.*, *J. Nat'l Cancer Inst.*, 92:1388-1402, 2000.

Therapeutically effective amount of a protein: A quantity of a protein sufficient to achieve a desired effect in a subject being treated. For instance, when
30 referring to a functionalized TGF- β family protein fusion this can be the amount necessary to induce a dose-dependent tissue specific effect. Examples of clinically

relevant TGF- β functionality include:

- the amount of TGF- β that, when applied topically to mucosal surfaces, can inhibit cycling of basal epithelial cells;
- the amount of TGF- β that, when administered topically to the epidermis, can enhance collagen deposition and promote accelerated wound healing; and
- the amount of TGF- β that can suppress local inflammatory reactions when administered locally at various sites.

An effective amount of a protein (such as a fusion protein of the disclosure) may be administered in a single dose, or in several doses, for example daily, during a course of treatment. However, the effective amount of protein will be dependent on the protein applied, the subject being treated, the severity and type of the affliction, and the manner of administration of the protein. For example, a therapeutically effective amount of a fusion protein can vary from about 0.01 mg/kg body weight to about 1 g/kg body weight.

The fusion proteins disclosed herein have equal application in medical and veterinary settings. Therefore, the general term "subject being treated" is understood to include all animals (*e.g.* humans, apes, dogs, cats, horses, and cows).

Transformed: A transformed cell is a cell into which has been introduced a nucleic acid molecule by molecular biology techniques. As used herein, the term transformation encompasses all techniques by which a nucleic acid molecule might be introduced into such a cell, including transfection with viral vectors, transformation with plasmid vectors, and introduction of naked DNA by electroporation, lipofection, and particle gun acceleration.

Vector: A nucleic acid molecule as introduced into a host cell, thereby producing a transformed host cell. A vector may include nucleic acid sequences that permit it to replicate in a host cell, such as an origin of replication. A vector may also include one or more selectable marker genes and other genetic elements known to those of ordinary skill in the art.

Unless otherwise explained, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art

to which this invention belongs. The singular terms "a," "an," and "the" include plural referents unless context clearly indicates otherwise. Similarly, the word "or" is intended to include "and" unless the context clearly indicates otherwise. It is further to be understood that all base sizes or amino acid sizes, and all molecular weight or molecular mass values, given for nucleic acids or polypeptides are approximate, and are provided for description. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the compositions and methods herein, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including explanations of terms, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

15 *III. Description of Several Embodiments*

This disclosure provides functionalized TGF- β fusion proteins that maintain substantial TGF- β biological activity. These fusion proteins are achieved by placing a functionalizing peptide between the pro- and active (mature) portions of a TGF- β protein, or at a relatively non-conserved site within the mature region of a TGF- β protein, or within a few residues (for instance within 15 or fewer amino acid residues) of the maturation cleavage site between the pro- and active (mature) portions of the TGF- β protein.

Encompassed herein are functional TGF- β family fusion proteins that contain a functionalizing peptide portion for detecting, quantifying or providing a specific additional function to the fusion protein and a mature TGF- β family protein, both as a monomer and in the form of a dimer (*e.g.*, a homodimer). Also encompassed are nucleic acid molecules encoding such fusion proteins, and conservative substitutions of such molecules.

Optionally, functionalized fusion proteins disclosed herein can also include a pro-region (latency associated peptide) of a TGF- β family protein, which for instance can be located to provide targeting and/or assembly and/or processing of

the fusion protein. In certain embodiments, the pro-region is located at the N-terminal region of the fusion protein.

Also encompassed herein are fusion proteins in which the TGF- β family member portion of the fusion protein is a functional variant of a naturally occurring
5 TGF- β family protein.

TGF- β family proteins useful for providing portions of the encompassed fusions proteins include TGF- β 2, TGF- β 3, TGF- β 1, TGF- β 4 (chicken), TGF- β 5 (*Xenopus*), GDF-9 (mouse/human), BMP-16/nodal (mouse), Fugacin (*Xenopus*), BMP3, Sumitomo-BIP/GDF-10 (mouse), ADMP (*Xenopus*), BMP-9, Dorsalin-1
10 (Chicken), BMP-10, BMP-13/GDF-6 (mouse), Radar (Zebrafish), GDF-1/CDMP-1 (mouse/human), BMP-12/GDF-7 (mouse), BMP-5, BMP-6, BMP-7/OP-1, BMP-8/OP-2, PC8/OP-3 (mouse), 60A (*Drosophila*), BMP-2, BMP-4, Decapentaplegic (*Drosophila*), Vg-1 (*Xenopus*), Univin (sea urchin), Vgr-2/GDF-3, GDF-1, Screw (*Drosophila*), BMP-11, GDF-8, Activin β C, Activin β D (*Xenopus*), Activin β E, BMP-
15 14/GDF-12, Activin β A, Activin β B, GDF-14, Mullerian inhibiting substance, and α -inhibin. Specific examples of disclosed fusions contain mature and/or pro-regions from TGF- β 1, TGF- β 2, or TGF- β 3. The mature and pro-region contained with a single fusion are not necessarily derived from the same TGF- β family protein, though in some embodiments the mature and pro-regions will be derived from the
20 same native TGF- β family protein.

Dimers formed from the functional TGF- β family fusion proteins can be made, for instance, by methods that include expressing a nucleic acid molecule in a eukaryotic cell to produce a monomer fusion protein. The monomers can then be associated together to form a dimer. Such nucleic acid molecules can include a
25 sequence encoding a functionalizing peptide portion of the functional TGF- β family fusion protein, a sequence encoding the desired mature TGF- β family protein, and a sequence encoding a pro-region (latency associated peptide) of a TGF- β family protein. In certain embodiments, the pro-region (latency associated peptide) is located to provide targeting and/or assembly and/or processing of the fusion protein
30 encoded for by the nucleic acid. This will often be at or near the N-terminal end of

the overall fusion protein construct. Thus, the sequence encoding the pro-region may be located upstream to both the sequence encoding the functionalizing peptide portion and the sequence encoding the mature TGF- β family protein.

In certain embodiments, the pro-region (latency associated peptide) is
5 cleaved from the functionalized TGF- β fusion protein monomers disclosed herein. Likewise, in certain embodiments the pro-region is cleaved from one or both of the monomers in dimers disclosed herein. Methods are provided for actively cleaving the pro-region from the monomers, for instance through the addition of a protease to a sample of the monomer. Alternatively, the pro-region in some instances is
10 removed through the expression of the monomer construct in a cell or system that contains such a protease.

In specific embodiments, the functionalizing peptide portion of a disclosed TGF- β fusion protein is at the N-terminus of the mature TGF- β family protein portion of the fusion. Such a fusion protein is exemplified by NFLAG-TGF- β 1;
15 nucleic acid and/or amino acid sequences related to this fusion are shown in SEQ ID NOs: 8-11.

In other specific embodiments, the functionalizing peptide portion is inserted within the mature functional TGF- β family protein. Such insertion can be, for instance, at a position of relatively low sequence conservation within the TGF- β
20 super family. In examples of such fusion proteins, the functionalizing peptide portion is inserted between about residues 11 and 12 of the mature TGF- β family protein. Such a fusion protein is exemplified by 11/12FLAG-TGF- β 1; nucleic acid and/or amino acid sequences related to this fusion are shown in SEQ ID NOs: 12-15.

Sub B3 In other examples of fusion proteins in which the functionalizing peptide is
25 inserted within the mature TGF- β family protein, a portion of the TGF- β family protein may be repeated both before and after the inserted peptide. For instance, in some specific examples, the peptide is inserted after five amino acid residues of the mature TGF- β family protein, and these five amino acids are then repeated after the peptide (such that the entire TGF- β mature protein occurs in the fusion after the
30 peptide). Such a fusion protein is exemplified by N+5FLAG-TGF- β 1 (SEQ ID

2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500 2501 2502 2503 2504 2505 2506 2507 2508 2509 2510 2511 2512 2513 2514 2515 2516 2517 2518 2519 2520 2521 2522 2523 2524 2525 2526 2527 2528 2529 2530 2531 2532 2533 2534 2535 2536 2537 2538 2539 2540 2541 2542 2543 2544 2545 2546 2547 2548 2549 2550 2551 2552 2553 2554 2555 2556 2557 2558 2559 2560 2561 2562 2563 2564 2565 2566 2567 2568 2569 2570 2571 2572 2573 2574 2575 2576 2577 2578 2579 2580 2581 2582 2583 2584 2585 2586 2587 2588 2589 2590 2591 2592 2593 2594 2595 2596 2597 2598 2599 2600 2601 2602 2603 2604 2605 2606 2607 2608 2609 2610 2611 2612 2613 2614 2615 2616 2617 2618 2619 2620 2621 2622 2623 2624 2625 2626 2627 2628 2629 2630 2631 2632 2633 2634 2635 2636 2637 2638 2639 2640 2641 2642 2643 2644 2645 2646 2647 2648 2649 2650 2651 2652 2653 2654 2655 2656 2657 2658 2659 2660 2661 2662 2663 2664 2665 2666 2667 2668 2669 2670 2671 2672 2673 2674 2675 2676 2677 2678 2679 2680 2681 2682 2683 2684 2685 2686 2687 2688 2689 2690 2691 2692 2693 2694 2695 2696 2697 2698 2699 2700 2701 2702 2703 2704 2705 2706 2707 2708 2709 2710 2711 2712 2713 2714 2715 2716 2717 2718 2719 2720 2721 2722 2723 2724 2725 2726 2727 2728 2729 2730 2731 2732 2733 2734 2735 2736 2737 2738 2739 2740 2741 2742 2743 2744 2745 2746 2747 2748 2749 2750 2751 2752 2753 2754 2755 2756 2757 2758 2759 2760 2761 2762 2763 2764 2765 2766 2767 2768 2769 2770 2771 2772 2773 2774 2775 2776 2777 2778 2779 2780 2781 2782 2783 2784 2785 2786 2787 2788 2789 2790 2791 2792 2793 2794 2795 2796 2797 2798 2799 2800 2801 2802 2803 2804 2805 2806 2807 2808 2809 2810 2811 2812 2813 2814 2815 2816 2817 2818

15

25

30

Further embodiments are methods of adding a non-native functionality to a mature biologically active TGF- β family protein. Such methods involve inserting a functionalizing peptide portion (such as a tag, a targeting moiety, or a biologically active protein domain) between a TGF- β pro-region and a TGF- β mature protein, or
5 at a relatively non-conserved site within the mature region of a TGF- β family protein. Representative TGF- β family proteins that can be used to construct such fusions include TGF- β 2, TGF- β 3, TGF- β 1, TGF- β 4 (chicken), TGF- β 5 (*Xenopus*), GDF-9 (mouse/human), BMP-16/nodal (mouse), Fugacin (*Xenopus*), BMP3, Sumitomo-BIP/GDF-10 (mouse), ADMP (*Xenopus*), BMP-9, Dorsalin-1 (Chicken),
10 BMP-10, BMP-13/GDF-6 (mouse), Radar (Zebrafish), GDF-1/CDMP-1 (mouse/human), BMP-12/GDF-7 (mouse), BMP-5, BMP-6, BMP-7/OP-1, BMP-8/OP-2, PC8/OP-3 (mouse), 60A (*Drosophila*), BMP-2, BMP-4, Decapentaplegic (*Drosophila*), Vg-1 (*Xenopus*), Univin (sea urchin), Vgr-2/GDF-3, GDF-1, Screw (*Drosophila*), BMP-11, GDF-8, Activin β C, Activin β D (*Xenopus*), Activin β E, BMP-
15 14/GDF-12, Activin β A, Activin β B, GDF-14, Mullerian inhibiting substance, and α -inhibin.

Further embodiments include methods using TGF- β fusion proteins disclosed herein to treat a disease that responds to administration of a TGF- β family protein, or assess a pharmacologic property of the protein. Such methods involve
20 administering a therapeutically sufficient amount of a disclosed fusion protein to a subject. In certain of such methods, the functionalizing peptide portion of the fusion is for detection or quantifying the fusion protein, and specific examples of such methods further involve detecting and/or quantifying the fusion protein. A pharmacokinetic or pharmacodynamic calculation can be performed on data
25 generated using such methods. Such calculation can involve a pharmacodynamic calculation of cellular site of protein action, dose-response relationship(s), structure-activity relationship(s), or quantitation of protein-receptor interaction(s). Alternatively, such calculations can involve a pharmacokinetic calculation of absorption, bioavailability, distribution, metabolism, or elimination/clearance of the
30 protein.

IV. *Functionalized TGF- β Family Protein Fusions*

The current disclosure provides TGF- β fusion proteins that can be expressed in mammalian cells while maintaining physiological function(s) of the fusion, and in particular the TGF- β family functionality of the fusion protein. The TGF- β family protein is genetically fused with a functionalizing peptide, such as a peptide tag or a targeting or other functional domain. Representative molecules constructed using the disclosed method comprise an epitope tag (such as the FLAG tag) fused to TGF- β 1 near the amino-terminus of the mature protein. This placement of the functionalizing peptide places it effectively in the middle of the TGF- β precursor protein, between the pro-region and the mature region or near this juncture. The tagged protein can then be expressed with the pro-region attached to enable proper folding of the TGF- β homodimer.

Tagging or otherwise generating a fusion of a TGF- β protein is complicated by the fact that proteins in the TGF- β family are initially synthesized as inactive, latent precursors (Roberts, *Annu. Rev. Immunol.* 16:137-61:137-161, 1998) that contain a relatively large N-terminal pro-region, also referred to as the latency associated peptide (LAP). LAP is required for both proper folding and secretion of TGF- β family proteins. Removal of LAP from the mature region prior to expression results in a loss of biological activity. Previous attempts to add epitope tags to TGF- β 1 resulted in biologically inactive fusions (Wakefield *et al. Growth Factors*, 5:243-253, 1991). The TGF- β family fusion proteins described herein overcome the difficulties encountered by previous constructs, particularly low bio-activity and poor folding, because the functionalizing peptide portion of the fusion is placed at the pro-mature junction of the TGF- β family protein, or at a relatively unconserved position within the mature portion of the protein. This permits the LAP to provide folding, assembly, and targeting activity while still producing a mature fusion with the functionalizing peptide intact.

Two representative functionalized TGF- β fusion proteins described herein are FLAG-TGF- β 1 (wherein the FLAG epitope tag is inserted immediately before the N-terminus of mature TGF- β 1) and 11/12FLAG (wherein the FLAG epitope tag

is inserted between amino acid residues 11 and 12 of the mature TGF- β 1). Each of these fusions is demonstrated to be functional, based on *in vitro* and *in vivo* biological and biochemical analyses. Although the amount of FLAG-tagged molecules secreted was lower than un-tagged control TGF- β 1, even dilutions of the
5 FLAG-TGF- β 1 culture supernatants as great as 1:256 were still as effective as supernatants from cells transfected with WT TGF- β 1 in inhibiting cell growth. This indicates that the functionalized TGF- β fusions may have higher specific activity than does native TGF- β .

With the provision herein of functionalized TGF- β fusion proteins, and in particular the description of specific fusions, the construction of other TGF- β
10 fusions that maintain native bio-activity while having an addition function conveyed by a functionalizing peptide (such as a tag, a targeting moiety, or a biologically active protein), is now enabled. The construction of such additional molecules will now be described.

15 **A. Selection of fusion components.**

The choice of appropriate protein domains and/or peptides for incorporation into the disclosed functionalized TGF- β fusion proteins will be dictated by the desired specificity (or lack of specificity) of the TGF- β portion and the desired added function (*e.g.*, identification or isolation of the fusion protein, targeting of the
20 fusion, and so forth).

Fusion proteins of this disclosure include at least a functional portion of a TGF- β family protein. Representative members of the TGF- β family of proteins include: TGF- β 2, TGF- β 3, TGF- β 1, TGF- β 4 (chicken), TGF- β 5 (*Xenopus*), GDF-9 (mouse/human), BMP-16/nodal (mouse), Fugacin (*Xenopus*), BMP3, Sumitomo-
25 BIP/GDF-10 (mouse), ADMP (*Xenopus*), BMP-9, Dorsalin-1 (Chicken), BMP-10, BMP-13/GDF-6 (mouse), Radar (Zebrafish), GDF-1/CDMP-1 (mouse/human), BMP-12/GDF-7 (mouse), BMP-5, BMP-6, BMP-7/OP-1, BMP-8/OP-2, PC8/OP-3 (mouse), 60A (*Drosophila*), BMP-2, BMP-4, Decapentaplegic (*Drosophila*), Vg-1 (*Xenopus*), Univin (sea urchin), Vgr-2/GDF-3, GDF-1, Screw (*Drosophila*), BMP-
30 11, GDF-8, Activin β C, Activin β D (*Xenopus*), Activin β E, BMP-14/GDF-12,

Activin β A, Activin β B, GDF-14, Mullerian inhibiting substance, and α -inhibin. Any of the listed TGF- β family proteins, and any other proteins identified as belonging to the TGF- β family (or "superfamily") can be used to construct fusions. Different members of the family can be selected depending on, for instance, the biological system in which the fusion is intended to function (*e.g.*, mammalian) and/or the desired biological specificity of TGF- β activity. Specificities of the listed TGF- β family proteins are known to one of ordinary skill in the art. See, for instance, Doetschman, *Lab. Anim. Sci.* 49:137-143, 1999; Letterio and Roberts, *Annu. Rev. Immunol.* 16:137-61:137-161, 1998; Wahl, *J. Exp. Med.* 180:1587-1590, 1994; Letterio, Roberts, *J. Leukoc. Biol.* 59:769-774, 1996; Piek *et al.*, *FASEB J.* 13:2105-2124, 1999; Heldin *et al.*, *Nature* 390:465-471, 1979; and De Caestecker *et al.*, *J. Nat'l Cancer Inst.*, 92:1388-1402, 2000. Also, see Roberts and Sporn, *Mol. Reprod. Dev.* 32:91-98, 1992; Roberts *et al.*, *Ciba. Found. Symp.* 157:7-15, 1991; Kingsley, *Genes Dev.* 8:133-146, 1994; Roberts and Sporn, "The transforming growth factors- β ." In: *Peptide Growth Factors and their Receptors*. (Sporn and Roberts, eds.), 95:419-472, 1990.

The pro-region of a TGF- β family protein can be included in the functionalized fusion protein. Such inclusion is advantageous for the correct folding, dimerization, and secretion of the fusion protein in eukaryotic cell systems, and may in some instances be necessary to achieve maximal TGF- β biological activity in the resultant fusion. The pro-region is usually placed upstream (towards the amino-terminus) of the mature portion of the TGF- β protein, and may be cleaved off during production of the protein. In most instances, removal of the pro-region is necessary for complete activity of the fusion protein, since it is effective at blocking binding of the TGF- β protein at its receptor(s). Though the detailed examples herein describe constructs in which the pro-region and mature regions are selected from the same TGF- β protein, it is believed that this is not necessary for the function of the disclosed fusion proteins.

Likewise, selection of a functionalizing peptide portion for a fusion protein of the current disclosure will depend on the purpose of the fusion. Functionalizing peptide portions can be short peptides (such as tags), or longer peptide molecules or

protein domains or proteins (such as targeting domains or enzymatically active proteins). Merely by way of example, functionalizing peptide tags that can be used in fusions according to this disclosure include epitope tags (such as myc, T7, GST, HA, or FLAG), translocation/transduction tags (such as Tat), purification tags (such as the hexa-histidine tag) and other peptide labels, such as green fluorescent protein (GFP). An epitope tag can be added to a TGF- β fusion protein in order to allow the resultant fusion protein to be identified through the use of an antibody that recognizes the epitope. A purification tag can be added to a TGF- β fusion protein in order to allow the resultant fusion protein to be purified, for instance through column chromatography.

The fusion proteins as provided herein also can be functionalized through the addition of a cell-targeting protein domain or protein. Such domains can be used to specifically target the TGF- β fusion to a specific cell type, tissue type, or organ (the "target" of that fusion protein). Targeting of TGF- β fusion proteins to specific tissues (cells, organs, etc.) may reduce undesirable side effects of systemic administration of such proteins, including the activity of the fusion proteins in a spectrum of tissues throughout the subject. Cell-targeting domains/proteins include, but are not limited to proteins (or domains of proteins) that bind to specific cell-surface receptors, including receptors found on only specific cell types, at specific developmental stages, under specific environmental or clinical conditions, or in particular disease states. Given the ability of TGF- β to inhibit T-cells, antigens that are specifically associated with T-cell activation might be particularly useful, including CD40 ligand and CTLA4. Each are expressed at high levels on activated T-cells, but not on the resting T-cell. More importantly the receptor for CD40 and ligand for CTLA4 have been defined and could potentially be used to target TGF- β fusion proteins to T cells. The selection of a targeting portion of a TGF- β fusion protein will be influenced by the desired target (cell, tissue, organ, etc.) to which the fusion is intended to bind; one of ordinary skill in the relevant art will know which targeting portions are appropriate for directing the fusion to specific targets. Targeting portions of the subject fusions can serve additional purposes (provide additional functions) within the same fusion protein, for instance identification (*e.g.*,

through an epitope of the targeting portion, or an increase in molecular weight) and/or purification of the fusion.

The functionalizing portion of fusions also can be a protein or domain that has a function independent of tagging or targeting, for instance an enzymatic, catalytic, or other biological function, or as a stabilizing influence that renders the fusion less prone to proteolysis or other removal from the system.

The choice of appropriate linker, if any linker is used, also will be influenced by the function of the two portions of the molecule, and whether these two portions can or must interact or should or can be held apart from each other. In general, a linker used in a functionalized TGF- β fusion will be of a length and secondary character sufficient to permit the functionalizing portion of the fusion to perform its function without hindering the activity of the TGF- β portion. Linkers can be as simple as a few amino acids that are included to facilitate construction of the fusion, for instance by the addition of one or more restriction endonuclease sites in the corresponding recombinant nucleic acid fusion molecule.

B. Assembly.

The construction of fusion proteins from domains of known proteins, or from whole proteins or proteins and peptides, is well known. In general, a nucleic acid molecule that encodes the desired protein and/or peptide portions are joined using genetic engineering techniques to create a single, operably linked fusion oligonucleotide. Appropriate molecular biological techniques may be found in Sambrook *et al.* (1989). Examples of genetically engineered multi-domain proteins, including those joined by various linkers, and those containing peptide tags, can be found in the following patent documents:

- U.S. Patent No. 5,994,104 ("Interleukin-12 fusion protein");
- U.S. Patent No. 5,981,177 ("Protein fusion method and construction");
- U.S. Patent No. 5,914,254 ("Expression of fusion polypeptides transported out of the cytoplasm without leader sequences");
- U.S. Patent No. 5,856,456 ("Linker for linked fusion polypeptides");
- U.S. Patent No. 5,767,260 ("Antigen-binding fusion proteins");
- U.S. Patent No. 5,696,237 ("Recombinant antibody-toxin fusion protein");

U.S. Patent No. 5,587,455 ("Cytotoxic agent against specific virus infection");

U.S. Patent No. 4,851,341 ("Immunoaffinity purification system");

U.S. Patent No. 4,703,004 ("Synthesis of protein with an identification peptide"); and

WO 98/36087 ("Immunological tolerance to HIV epitopes").

The placement of the functionalizing peptide portion within the subject fusion proteins is influenced by the activity of the functionalizing peptide portion and the need to maintain at least substantial TGF- β biological activity in the fusion. Two methods for placement of a functionalizing peptide are illustrated in the detailed examples: between the pro-region and the mature region of the TGF- β family protein portion of the fusion, and at a location within the mature TGF- β family protein portion that exhibits either low sequence conservation or amenability to insertions. Though these are not the only locations in which functionalizing peptides can be inserted, they serve as good examples, and will be used as illustrations. Other appropriate insertion locations can be identified by inserting test peptide encoding sequences (*e.g.*, a sequence encoding the FLAG peptide) into a TGF- β construct at different locations, then assaying the resultant fusion for TGF- β biological activity and functionalizing peptide activity, using assays that are appropriate for the specific portions used to construct the fusion. The activity of TGF- β family proteins can be measured using any of various known techniques, including those method described herein (see, *e.g.*, section V, below, and Example 2).

C. Expression.

One skilled in the art will understand that there are myriad ways to express a recombinant protein such that it can subsequently be purified. In general, an expression vector carrying the nucleic acid sequence that encodes the desired protein will be transformed into a microorganism for expression. Such microorganisms can be prokaryotic (bacteria) or eukaryotic (*e.g.*, yeast). One appropriate species of bacteria is *Escherichia coli* (*E. coli*), which has been used extensively as a

laboratory experimental expression system. A eukaryotic expression system will be preferred where the protein of interest requires eukaryote-specific post-translational modifications such as glycosylation. Also, protein can be expressed using a viral (*e.g.*, vaccinia) based expression system.

5 Protein can also be expressed in animal cell tissue culture, and such a system will be appropriate where animal-specific protein modifications are desirable or required in the recombinant protein. Such expression is particularly appropriate where native assembly and export of the TGF- β fusion protein is desirable, since the activity of TGF- β family proteins is influenced by native dimerization (folding and
10 assembly) and secretion from the cell.

The expression vector can include a sequence encoding a targeting peptide positioned in such a way as to be fused to the coding sequence of the functionalized TGF- β fusion protein. One such "targeting" peptide is the precursor region of the TGF- β protein itself, which is important in native export of the TGF- β family
15 protein from the cell. Targeting peptides/domains included in the fusion can be in addition to another functionalizing peptide (such as a tag). Targeting domains may allow the fusion protein to be targeted to specific extra-cellular locations, or simply to be secreted from the cell, and may be removed during or soon after synthesis of the fusion protein. In addition, multiple targeting peptides can be included in a
20 single fusion, for instance a peptide/domain that directs the fusion protein to be secreted, and another peptide/domain that directs the secreted protein to a target (cell, tissue, organ, etc.). Various appropriate prokaryotic and eukaryotic targeting peptides, and nucleic acid molecules encoding such, are known to one of ordinary skill in the art. Through the use of a eukaryotic secretion-type signal sequence, the
25 functionalized TGF- β fusion protein can be expressed in a transgenic animal (for instance a cow, pig, or sheep) in such a manner that the fusion protein is secreted into the milk of the animal. Targeting protein portions also may be used to ensure that a transgenically expressed fusion protein is secreted into the circulatory system of the transgenic animal, thereby permitting the fusion protein to be transported to a
30 target (cell, tissue, organ, etc.).

Vectors suitable for stable transformation of culturable cells are also well known. Typically, such vectors include a multiple-cloning site suitable for inserting a cloned nucleic acid molecule, such that it will be under the transcriptional control of 5' and 3' regulatory sequences. In addition, transformation vectors include one or
5 more selectable markers; for bacterial transformation this is often an antibiotic resistance gene. Such transformation vectors typically also contain a promoter regulatory region (*e.g.*, a regulatory region controlling inducible or constitutive expression), a transcription initiation start site, a ribosome binding site, an RNA processing signal, and a transcription termination site, each functionally arranged in
10 relation to the multiple-cloning site. For production of large amounts of recombinant proteins, an inducible promoter is preferred. This permits selective production of the recombinant protein, and allows both higher levels of production than constitutive promoters, and enables the production of recombinant proteins that may be toxic to the expressing cell if expressed constitutively.

15 In addition to these general guidelines, protein expression/purification kits are produced commercially. See, for instance, the QIAexpress™ expression system from QIAGEN (Chatsworth, CA) and various expression systems provided by INVITROGEN (Carlsbad, CA). Depending on the details provided by the
manufacturers, such kits can be used for production and purification of the disclosed
20 functionalized TGF-β fusion proteins.

D. Purification.

One skilled in the art will understand that there are myriad ways to purify recombinant polypeptides, and such typical methods of protein purification may be used to purify the disclosed functionalized TGF-β fusion proteins. Such methods
25 include, for instance, protein chromatographic methods including ion exchange, gel filtration, HPLC, monoclonal antibody affinity chromatography and isolation of insoluble protein inclusion bodies after over production. In addition, purification affinity-tags, for instance a six-histidine sequence, may be recombinantly fused to the protein and used to facilitate polypeptide purification (*e.g.*, in addition to another
30 functionalizing portion of the fusion, such as a targeting domain or another tag). A specific proteolytic site, for instance a thrombin-specific digestion site, can be

engineered into the protein between the tag and the remainder of the fusion to facilitate removal of the tag after purification, if such removal is desired.

Commercially produced protein expression/purification kits provide tailored protocols for the purification of proteins made using each system. See, for instance, the *QIAexpress*TM expression system from QIAGEN (Chatsworth, CA) and various expression systems provided by INVITROGEN (Carlsbad, CA). Where a commercial kit is employed to produce a functionalized TGF- β fusion protein, the manufacturer's purification protocol is a preferred protocol for purification of that protein. For instance, proteins expressed with an amino-terminal hexa-histidine tag can be purified by binding to nickel-nitrilotriacetic acid (Ni-NTA) metal affinity chromatography matrix (*The QIAexpressionist*, QIAGEN, 1997).

More generally, the binding specificities of either the TGF- β or a functionalizing peptide domain, or both, of the disclosed fusion protein may be exploited to facilitate specific purification of the proteins. A preferred method of performing such specific purification would be column chromatography using column resin to which a target molecule, or an appropriate epitope or fragment or domain of the target molecule, has been attached.

If the functionalized TGF- β fusion protein is produced in a secreted form, *e.g.* secreted into the milk of a transgenic animal, purification can be from the secreted fluid. Alternatively, purification may be unnecessary if it is appropriate to apply the fusion protein directly to the subject in the secreted fluid (*e.g.* milk).

V. Variation of Functionalized TGF- β Fusion Protein(s)

A. Sequence Variants

Certain functional characteristics of the fusion proteins disclosed herein lie not in the precise amino acid sequence of the proteins, but rather in the three-dimensional structure inherent in the amino acid sequences encoded by the DNA sequences. It is possible to recreate the functional characteristics of the fusion proteins or protein domains by recreating the three-dimensional structure, without necessarily recreating the exact amino acid sequence. This can be achieved by designing a nucleic acid sequence that encodes for the three-dimensional structure,

but which differs, for instance by reason of the redundancy of the genetic code. Similarly, the DNA sequence may also be varied, while still producing a functionalized TGF- β fusion protein. Such substitutions, however, may not produce functional variants of the disclosed fusion proteins if the substitutions are made at essential amino acid positions, for instance, binding-specificity essential residues within TGF- β family proteins (see, *e.g.*, Huang *et al.*, *J. Biol. Chem.*, 274:27754-27758, 1999), epitope-structural residues of functionalizing epitope tags, or targeting or enzymatically essential residues within other functionalizing protein portions. Thus, it is useful to assay the activity of variant fusion proteins (or the appropriate portion of the variant fusion proteins) using available protocols, including for instance those described herein.

Variant TGF- β fusion proteins include proteins that differ in amino acid sequence from the disclosed sequences, and sequence constructed from the disclosed protein portions, but that share structurally significant sequence homology with such proteins. Variation can occur in any single domain of the fusion protein (*e.g.* the functionalizing domain, the TGF- β family protein domain, or the linker if such is present in the fusion). Variation can also occur in more than one of such domains in any particular variant protein. Such variants may be produced by manipulating the nucleotide sequence of the, for instance a FLAG-TGF- β -encoding sequence, using standard procedures, including site-directed mutagenesis or mutagenic nucleic acid amplification (*e.g.*, using PCR). The simplest modifications involve the substitution of one or more amino acids for amino acids having similar biochemical properties. These so-called "conservative" substitutions are likely to have minimal impact on the activity of the resultant protein, especially when made outside of the binding site of each domain. Table 1 shows amino acids that may be substituted for an original or native amino acid in a protein, and which are regarded as conservative substitutions.

Table 1

	Original Residue	Conservative Substitutions
	Ala	ser
	Arg	lys
5	Asn	gln; his
	Asp	glu
	Cys	ser
	Gln	asn
	Glu	asp
10	Gly	pro
	His	asn; gln
	Ile	leu; val
	Leu	ile; val
	Lys	arg; gln; glu
15	Met	leu; ile
	Phe	met; leu; tyr
	Ser	thr
	Thr	ser
	Trp	tyr
20	Tyr	trp; phe
	Val	ile; leu

More substantial changes in protein structure may be obtained by selecting one or more amino acid substitutions that are less conservative than those listed in Table 1. Such changes include changing residues that differ more significantly in their effect on maintaining polypeptide backbone structure (*e.g.*, sheet or helical conformation) near the substitution, charge or hydrophobicity of the molecule at the target site, or bulk of a specific side chain. The following substitutions are generally expected to produce the greatest changes in protein properties: (a) a hydrophilic residue (*e.g.*, seryl or threonyl) is substituted for (or by) a hydrophobic residue (*e.g.*, leucyl, isoleucyl, phenylalanyl, valyl or alanyl); (b) a cysteine or proline is substituted for (or by) any other residue; (c) a residue having an electropositive side chain (*e.g.*, lysyl, arginyl, or histadyl) is substituted for (or by) an electronegative residue (*e.g.*, glutamyl or aspartyl); or (d) a residue having a bulky side chain (*e.g.*, phenylalanine) is substituted for (or by) one lacking a side chain (*e.g.*, glycine).

Variant TGF- β domain or fusion protein-encoding sequences may be produced by standard DNA mutagenesis techniques, for example, M13 primer mutagenesis. Details of these techniques are provided in Sambrook (Ch. 15, In

Molecular Cloning: A Laboratory Manual, CSHL, New York, 1989). By the use of such techniques, variants may be created which differ in minor ways from native TGF- β encoding sequences (such as those encoding the proteins listed in FIG 1). DNA molecules and nucleotide sequences which are derivatives of native TGF- β encoding sequences and that differ from such sequence by the deletion, addition, or substitution of nucleotides while still encoding a protein that has TGF- β biological activity (either of the same specificity as the original TGF- β family member, or the specificity of another family member), are comprehended by this disclosure. In their most simple form, such variants may differ from the disclosed sequences by alteration of the coding region to fit the codon usage bias of the particular organism into which the molecule is to be introduced.

Alternatively, the coding region may be altered by taking advantage of the degeneracy of the genetic code to alter the coding sequence such that, while the nucleotide sequence is substantially altered, it nevertheless encodes a protein having an amino acid sequence substantially similar to the disclosed fusion sequences. For example, because of the degeneracy of the genetic code, four nucleotide codon triplets – (GCT, GCG, GCC and GCA) - code for alanine. The coding sequence of any specific alanine residue within a subject fusion protein, therefore, could be changed to any of these alternative codons without affecting the amino acid composition or characteristics of the encoded protein. Based upon the degeneracy of the genetic code, variant DNA molecules may be derived from the cDNA and gene sequences disclosed herein using standard DNA mutagenesis techniques as described above, or by synthesis of DNA sequences. Thus, this disclosure also encompasses nucleic acid sequences which encode a neutralizing bispecific fusion protein, but which vary from the disclosed nucleic acid sequences by virtue of the degeneracy of the genetic code.

Some variant TGF- β superfamily proteins previously have been identified or generated (see, *e.g.*, U.S. Patent No. 4,886,747; Yamada *et al.*, *J. Bone Miner. Res.*, 15:415-420, 2000; Huang *et al.*, *J. Biol. Chem.*, 274:27754-27758, 1999; Pociot *et al.*, *J. Am. Soc. Nephrol.*, 9:2302-2307, 1998; and Wharton *et al.*, *Genetics*, 142:493-505, 1996).

B. Peptide Modifications

The present disclosure includes biologically active molecules that mimic the action of the functionalized TGF- β fusion proteins provided herein, and specifically that maintain a TGF- β family protein activity and the desired additional

5 functionality (*e.g.*, targeting, recognition by a specific antibody (through an epitope), ease of purification, enzymatic functionality, and so forth) depending on the functionalizing portion that is fused to the TGF- β portion. The proteins of the disclosure include synthetic embodiments of fusion proteins described herein, as well as analogues molecules (non-peptide organic molecules), derivatives
10 (chemically functionalized protein molecules obtained starting with the disclosed peptide sequences) and variants (homologs) of these proteins that specifically maintain at least one TGF- β specific biological activity (*e.g.*, binding to a specific TGF- β receptor) and at least one additional function provided by the “functionalizing” portion of the fusion (which function will be dependent on the
15 portion chosen). Each protein of the disclosure is comprised of a sequence of amino acids, which may be either L- and/or D- amino acids, naturally occurring and otherwise.

Proteins may be modified by a variety of chemical techniques to produce derivatives having essentially the same activity as the unmodified proteins, and
20 optionally having other desirable properties. For example, carboxylic acid groups of the protein, whether carboxyl-terminal or side chain, may be provided in the form of a salt of a pharmaceutically-acceptable cation or esterified to form a C₁-C₁₆ ester, or converted to an amide of formula NR₁R₂ wherein R₁ and R₂ are each independently H or C₁-C₁₆ alkyl, or combined to form a heterocyclic ring, such as a 5- or 6-
25 membered ring. Amino groups of the protein, whether amino-terminal or side chain, may be in the form of a pharmaceutically-acceptable acid addition salt, such as the HCl, HBr, acetic, benzoic, toluene sulfonic, maleic, tartaric and other organic salts, or may be modified to C₁-C₁₆ alkyl or dialkyl amino or further converted to an amide.

30 Hydroxyl groups of the protein side chains may be converted to C₁-C₁₆ alkoxy or to a C₁-C₁₆ ester using well-recognized techniques. Phenyl and phenolic

rings of the protein side chains may be substituted with one or more halogen atoms, such as fluorine, chlorine, bromine or iodine, or with C₁-C₁₆ alkyl, C₁-C₁₆ alkoxy, carboxylic acids and esters thereof, or amides of such carboxylic acids. Methylene groups of the protein side chains can be extended to homologous C₂-C₄ alkenes.

5 Thiols can be protected with any one of a number of well-recognized protecting groups, such as acetamide groups. Those skilled in the art will also recognize methods for introducing cyclic structures into the proteins provided herein to select and provide conformational constraints to the structure that result in enhanced stability.

10 Peptidomimetic and organomimetic embodiments are also within the scope of the present disclosure, whereby the three-dimensional arrangement of the chemical constituents of such peptido- and organomimetics mimic the three-dimensional arrangement of the protein backbone and component amino acid side chains in a functionalized TGF- β fusion protein, resulting in such peptido- and
15 organomimetics of the proteins of this disclosure having at least one TGF- β biological activity and an additional function (conveyed by the "functionalizing" portion of the molecule, such as a tag or targeting portion, as discussed herein). For computer modeling applications, a pharmacophore is an idealized, three-dimensional definition of the structural requirements for biological activity. Peptido- and
20 organomimetics can be designed to fit each pharmacophore with current computer modeling software (using computer assisted drug design or CADD). See Walters, "Computer-Assisted Modeling of Drugs", in Klegerman & Groves, eds., 1993, *Pharmaceutical Biotechnology*, Interpharm Press: Buffalo Grove, IL, pp. 165-174 and *Principles of Pharmacology* Munson (ed.) 1995, Ch. 102, for descriptions of
25 techniques used in CADD. Also included within the scope of the disclosure are mimetics prepared using such techniques that produce biologically active, functionalized TGF- β fusion proteins.

C. Domain length variation.

It will be appreciated that the protein/peptide domains of the current
30 disclosure may be combined to produce fusion protein molecules without necessarily splicing the components in the same place. It is believed to be possible

to use shorter or longer fragments of each component domain. Such variation in length can be caused by the inclusion of longer or short functionalizing portions in the fusion, including for instance a linker portion. It is, however, in some instances not beneficial to use so short a portion of a tag or targeting peptide that the tag or
5 targeting peptide is no longer functional (and therefore cannot provide an additional function to the fusion protein).

D. Functionalizing peptide location in the fusion.

The placement of the functionalizing peptide portion within the subject fusion proteins is influenced by the activity of the functionalizing peptide portion
10 and the need to maintain at least substantial TGF- β biological activity in the fusion (as described herein). However, within these limitations, variants of the described functionalized TGF- β fusion molecules can be constructed by altering the placement of the functionalizing peptide. Such variants include moving the peptide insertion point a few (*e.g.*, one to about 10) amino acid residues towards the amino-or
15 carboxy-terminal end of the fusion protein (*e.g.*, inserting a FLAG or other tag between residues 12 and 13 instead of 11 and 12 of TGF- β 1, or between residues 21 and 22, or between residues 10 and 11, for instance). Likewise, a peptide tag could be inserted between any pair of adjacent residues between about residue 1 and about residue 22 of the mature portion of a TGF- β family protein. In some specific
20 examples, such insertion within the TGF- β family protein is accompanied by the removal of one or more of the TGF- β family protein amino acid residues. For instance, an peptide tag (such as the FLAG tag) could be inserted between residues 10 and 12, replacing residue 11 of the TGF- β family protein.

25 **VI. Activity of Functionalized TGF- β Family Fusion Proteins**

It is important to assess the chemical, physical and biological activity of the disclosed fusion proteins. Among other uses, such assays permit optimization of the domains chosen, optimization of the placement of the functionalizing portion within the fusion protein, optimization of the length and conformation of the linkers used to
30 connect portions of the fusion, and determination of the effect(s) of variant amino acid changes in the fusion proteins. Appropriate control molecules can be included

in each activity assay. Such controls molecules can include individual domains used to construct the fusion (*e.g.*, a part or all of the mature or precursor TGF- β family protein), composite domains expressed as separate molecules and mixed in the reaction, for instance in a 1:1 molar ratio, or fusions that include only one portion of the functionalized TGF- β fusion coupled to another protein or peptide (*e.g.*, a different tag on the same TGF- β protein, or the same tag on a different subject protein, either another member of the TGF- β family or another protein).

A. TGF- β family protein activity

The biological activity of a TGF- β family protein portion of a subject fusion protein can be assayed using any various different known TGF- β activity assays, including those described in detail herein. TGF- β biological activities that may be displayed by the subject fusion proteins include, but are not limited to, activities linked to the mature region of the protein (*e.g.*, binding to one or more cell surface receptors, mediation of a cellular response, etc.) and activities linked to the TGF- β pro-region (*e.g.*, secondary folding and tertiary assembly of the TGF- β homodimer, processing, secretion from the cell, etc.). Each of these functions can be measured in a fusion protein, either alone or in some instances in combination.

Binding of a TGF- β fusion protein can be measured in competitive binding assays (as described herein and elsewhere, *e.g.*, in Qian *et al.*, *J. Biol. Chem.*, 271:30656-30662, 1996; The activity of the mature region of a TGF- β protein (including the TGF- β fusions of this disclosure) also can be measured by assaying one or more downstream effects of the protein in a biological system. Such effects include phosphorylation of one or more smad proteins (such as smad2, as described below), and growth inhibition (*e.g.*, inhibition of CCL64 mink lung epithelial cells, as described below). Other assays for TGF- β activities are described, for instance, in Gray and Mason, *Science*, 247:1328-1330, 1990; Tuan *et al.*, *Conn. Tiss. Res.*, 34:1-9, 1996; Wakefield *et al.*, *Growth Factors*, 5:243-253, 1991; Danielpour and Roberts, *J. Immunol Methods* 180:265-272, 1995; Danielpour, *J Immunol Methods*. 158:17-25, 1993; Danielpour *et al.*, *Growth Factors* 2:61-71, 1989; Randall *et al.*,

Immunol. Methods 164:61-67, 1993; Danielpour *et al.*, *J. Cell Physiol.* 138:79-86, 1989.

The pro-region is important for correct folding, assembly, and secretion of a TGF- β protein; thus, activity of a pro-region that is included in a fusion protein of
5 this disclosure can be measured by assaying for these functions of the fusion protein. Pro-region activity has been examined previously (see, *e.g.*, Dubois *et al.*, *J. Biol. Chem.*, 270:10618-10624, 1995; Gray and Mason, *Science*, 247:1328-1330, 1990; Murphy-Ullrich and Poczatek, *Cytokine Growth Factor Rev.* 11:59-69, 2000; Barcellos-Hoff, *J Mammary Gland Biol. Neoplasia* 1:353-363, 1996; Hellmich *et al.*, *Metabolism* 49:353-359, 2000; Saharinen *et al.*, *Cytokine Growth Factor Rev.*,
10 10:99-117, 1999, and the methods described in these references are appropriate for measuring the activity of a pro-region in the subject fusions.

B. Activity of a functionalizing peptide

The biological activity of a functionalizing peptide that is fused to a TGF- β
15 protein portion to form a fusion can be assayed independently of the TGF- β biological activity(s) of the fusion. The appropriate assay(s) for measuring functionalizing peptide activity will be dictated largely by the functionalizing peptide.

The functionality of an epitope tag can be tested by detecting the fusion
20 protein using an antibody (or antibody derivative) known to bind to the epitope, for instance in an immunoblot ("western"), ELISA, or other assay system; such techniques are well known. Other identification tags can be tested for functionality based on their intended method of identification – *e.g.*, based on differential mobility or other added function. The functionality of a purification tag can be tested by
25 using it to purify the fusion protein, for instance using column chromatography or other conventional techniques.

The effective functionality of a targeting domain within a fusion protein can be tested by examining the targeting of the fusion protein in an experimental or clinical system. Such targeting can be examined using conventional techniques, for
30 instance fractionation, *in situ* hybridization, or through cell or tissue-specific

biological effects that result from the targeting of the fusion protein (*e.g.*, TGF- β mediated effects caused by the delivery of the fusion protein).

Other passenger proteins can be assayed based on the native or expected function of the passenger protein. Assays appropriate for any particular passenger
5 protein will be specific to that passenger, and will be known to those of ordinary skill in the art.

VII. Incorporation of Fusion Proteins into Pharmaceutical Compositions

Pharmaceutical compositions that comprise at least one functionalized TGF-
10 β fusion protein as described herein as an active ingredient will normally be formulated with an appropriate solid or liquid carrier, depending upon the particular mode of administration chosen. The pharmaceutically acceptable carriers and excipients useful with compositions in this disclosure are conventional. For instance, parenteral formulations usually comprise injectable fluids that are
15 pharmaceutically and physiologically acceptable fluid vehicles such as water, physiological saline, other balanced salt solutions, aqueous dextrose, glycerol or the like. Excipients that can be included are, for instance, other proteins, such as human serum albumin or plasma preparations. If desired, the pharmaceutical composition to be administered may also contain minor amounts of non-toxic auxiliary
20 substances, such as wetting or emulsifying agents, preservatives, and pH buffering agents and the like, for example sodium acetate or sorbitan monolaurate.

Other medicinal and pharmaceutical agents, for instance anti-proliferative agents, anti-infectives, or anti-cancer agents, also may be included.

The dosage form of the pharmaceutical composition will be determined by
25 the mode of administration chosen. For instance, in addition to injectable fluids, topical and oral formulations can be employed. Topical preparations can include eye drops, ointments, sprays and the like. Oral formulations may be liquid (*e.g.*, syrups, solutions or suspensions), or solid (*e.g.*, powders, pills, tablets, or capsules). For solid compositions, conventional non-toxic solid carriers can include
30 pharmaceutical grades of mannitol, lactose, starch, or magnesium stearate. Actual

methods of preparing such dosage forms are known, or will be apparent, to those of ordinary skill in the art.

The pharmaceutical compositions that comprise functionalized TGF- β fusion protein(s) can be formulated in unit dosage form, suitable for individual
5 administration of precise dosages. Possible unit dosages may contain, for instance, approximately 1.0 μ g to approximately 100 μ g of protein. The amount of active compound administered will be dependent on the subject being treated, the severity of the affliction, and the manner of administration, and is best left to the judgment of the prescribing clinician. Within these bounds, the formulation to be administered
10 will contain a quantity of the active component(s) in an amount effective to achieve the desired effect in the subject being treated.

VIII. Uses of Functionalized TGF- β Fusion Proteins

The functionalized TGF- β fusion proteins described herein can be used in a
15 number of systems in place of native TGF- β molecules, including: (1) detection of tagged ligand in transfected cells; (2) detection of cell surface expression of TGF- β receptor complexes by flow cytometry; and (3) measurement of cell surface levels of receptor complexes in non-radioactive cross-linking assays. The functionalized TGF- β fusion proteins are safer, faster and cheaper alternatives to the use of [125 I]
20 radiolabeled TGF- β molecules and expands the repertoire of techniques that can be used to look at TGF- β receptor expression levels.

Because the subject functionalized TGF- β fusion proteins can be distinguished from native TGF- β , a clinically/experimentally administered, functionalized TGF- β molecule can be traced within a subject or experimental
25 system. This permits determination of the distribution (cellular, tissue, etc.), half-life, elimination, and circulating levels of the administered protein, as differentiated from any endogenous TGF- β , permitting the clinician to more finely tune dosages and administration regimes.

The tagged protein also can be used to study TGF- β receptor expression
30 levels, for instance in different tissues or at different times (*e.g.*, after different

clinical or experimental treatments). In addition to being useful in prognosis and diagnosis of disease states (see below), this makes the fusion proteins useful for testing possible drug or other therapeutic treatments, or other regimens that might intentionally or unintentionally alter the expression level (or stability) of a TGF- β receptor.

In the clinic, trials employing a TGF- β family protein (such as TGF- β 1) to look at amelioration of disease symptoms in, for instance, arthritis, multiple sclerosis, colitis or in other diseases, can now use a functionalized TGF- β fusion protein (such as FLAG-TGF- β 1). This allows the clinician to follow bio-availability of the TGF- β family ligand and to accurately quantify the level that is exogenously administered.

The functionalized TGF- β fusion proteins provided herein may be administered to humans, or other animals on whose cells they are effective, in various manners such as topically, orally, intravenously, intramuscularly, intraperitoneally, intranasally, intradermally, intrathecally, and subcutaneously. The particular mode of administration and the dosage regimen will be selected by the attending clinician, taking into account the particulars of the case (*e.g.*, the subject, the disease, the disease state involved, and whether the treatment is prophylactic or post-infection). Treatment may involve daily or multi-daily doses of fusion protein(s) over a period of a few days to months, or even years.

If treatment is through the direct administration of cells expressing the functionalized TGF- β fusion protein to the subject, such cells (*e.g.* transgenic pluripotent or hematopoietic stem cells or B cells) may be administered at a dose of between about 10^6 and 10^{10} cells, on one or several occasions. The appropriate number of cells will depend on the patient, as well as the fusion protein and cells chosen to express the protein.

A general strategy for transferring genes into donor cells is disclosed in U.S. Patent No. 5,529,774. Generally, a gene encoding a protein having therapeutically desired effects is cloned into a viral expression vector, and that vector is then introduced into the target organism. The virus infects the cells, and produces the protein sequence *in vivo*, where it has its desired therapeutic effect. See, for example,

Zabner *et al.*, *Cell* 75:207-216, 1993. As an alternative to adding the sequences encoding the functionalized TGF- β fusion protein or a homologous protein to the DNA of a virus, it is also possible to introduce such a gene into the somatic DNA of infected or uninfected cells, by methods that are well known in the art (Sambrook *et al.* (In
5 *Molecular Cloning: A Laboratory Manual*, CSHL, New York, 1989). These methods can be used to introduce the herein disclosed fusion proteins to human cells to provide long-term production of the fusion protein.

Fusion proteins disclosed herein can be used for the detection of TGF- β receptor expression in tissues or cells of patients with cancer and immune disorders.
10 In this instance, the ability of the epitope-tagged ligand to bind to receptors on fresh specimens or fixed tissues can be coupled with quantitative analysis of cell-bound epitope tag. This can be accomplished either by direct visualization with immunohistochemical analyses, or through the use of Fluorescence Activated Cell Sorting (FACS) to detect the presence of a fluorochrome-antibody conjugate raised
15 specifically to the epitope tag.

The presence of an epitope tag in functional TGF- β fusion proteins will allow for accurate assessment of the delivery, distribution, and elimination following systemic, topical, or enteral administration of the fusion TGF- β protein. This is impossible to do with the native protein, as the production of TGF- β via the
20 endogenous gene cannot be distinguished from the administered exogenous protein. Thus, this fusion protein will provide a handle for an unambiguous assessment of the pharmacodynamics of TGF β in the setting of clinical trials. This is also an important consideration for any future gene therapy approaches based on the delivery of a TGF- β expression system. The use of a tagged TGF- β in such vectors
25 will allow one to follow production, both local and systemic, in a quantitative fashion, over time. Examples of specific applications include:

6651 -- topical therapy of cutaneous wounds for the acceleration of wound healing;

6652 -- topical application to oral mucosa to prevent chemotherapy-induced
30 mucositis;

6653 -- systemic administration of TGF- β to prevent chemotherapy -induced myelosuppression;

6654 -- aerosolized administration of TGF- β for the treatment of reactive airway diseases such as allergen-induced asthma;

5 6655 -- topical treatment of inflammatory dermatitis;

6656 -- intranasal administration of cDNA expression vector encoding FLAG-TGF- β for the prevention and / or treatment of colitis (Kitani, *et al.*, *J. Exp. Med.* 192:41-52, 2000).

10 IX. *Experimental Animal Systems*

The teaching of adding a functionalizing peptide or protein or peptide portion to a functional TGF- β family protein now makes it possible to engineer TGF- β family protein "knock-ins" to examine whether a specific TGF- β (*e.g.*, TGF- β 1) can rescue the phenotype of TGF- β specific "knock-out" mice (such as TGF- β 2 or TGF-
15 β 3 knock-outs) without the complications caused by cross reactivity of species-specific or isoform-specific antibodies.

The essential question that "knock-in" studies aim to address is whether there is substantial redundancy between the individual isoforms of the TGF- β family expressed in mammals. While they have a great deal of similarity at the amino acid
20 level, it is not clear whether one isoform can sufficiently replace the other *in vivo* or can compensate for the absence of another *in vivo*. To determine this, we can now replace each TGF- β locus with either the same isoform or with one of the immediate or distant family members with the epitope tag inserted. The tag allows identification of when and where the specific cytokine is expressed. More
25 importantly, we can determine whether (for example) TGF- β 2, when expressed where and when TGF- β 1 is normally expressed, can exert the appropriate functions necessary to maintain homeostasis.

X. *Diagnosis and Prognosis*

30 The development and/or progression of many pathologic conditions, including cancer and immune disorders, is often associated with loss of expression

of TGF- β family protein receptors. The functionalized fusion proteins of the disclosure provide tools for quantitative, real-time measurement of receptor expression in tissues and cells by virtue of the ability of these fusions to bind to TGF- β receptors, and be identified (*e.g.*, by the presence of an epitope tag or other
5 identifiable protein or peptide portion). These molecules can be used to diagnose and/or determine progression of diseases associated with receptor loss.

Specific examples of diseases or other pathologic conditions that can be diagnosed or prognosed using the subject fusions proteins include: cancer (see Reiss, *Microb. Infect.*, 1:1327-1347, 1999; De Caestecker *et al.*, *J. Nat'l Cancer*
10 *Inst.* 92:1388-1402, 2000; Kim *et al.*, *Cyto. Growth Factor Rev.* 11:159-169, 2000; and Taketo *et al.*, *Cyto. Growth Factor Rev.* 11:147-159, 2000); sound healing (Ashcroft and Roberts, *Cyto. Growth Factor Rev.* 11:125-133, 2000; Roberts and Sporn, "Transforming growth factor- β ." In: *The molecular and cellular biology of wound repair*. Clark (Ed.), New York: Plenum Press, p275-308, 1996; and Beck *et*
15 *al.*, *J. Clin. Invest.* 92:2841-2849; 1993); atherosclerosis (McCaffery, *Cyto. Growth Factor Rev.* 11:103-114, 2000); kidney disease (Sharma and McGowan, *Cyto. Growth Factor Rev.* 11:115-125, 2000); and hereditary hemorrhagic telangiectasia (Johnson *et al.*, *Nat. Genetics* 13:189-195, 1996; and McAllister, *et al.*, *Nat. Genetics* 8:345-351, 1994). More generally, see Blobe *et al.* (*NEJM* 342:1350-
20 1358, 2000).

In each of these situations, an appropriate functionalized (*e.g.*, epitope tagged) TGF- β fusion protein can be applied to a cell sample from a subject, and the amount of binding of the fusion protein to the target TGF- β receptor can be determined and compared to the amount found in a control sample. The level of
25 ligand binding to the receptor is indicative of the level of the receptor in that sample. An altered receptor level (such as a reduced receptor level) is indicative of the indicated associated disease or pathologic condition.

XI. Kits

30 Kits are provided that contain at least one functionalized TGF- β fusion protein, or a nucleic acid molecule (*e.g.*, a vector) that encodes such a protein, or

both, in one or more contains. The provided kits may also include written instructions. The instructions can provide calibration curves or charts to compare with the determined (*e.g.*, experimentally measured) values. Included are kits that can be used for diagnosis or prognosis of a disease or other condition associated
5 with a change (decrease or increase) in the level of cell surface expression of a TGF- β receptor.

The invention is further illustrated by the following non-limiting Examples.

10

EXAMPLES

Example 1: Construction and expression of FLAG-TGF- β 1 and 11/12FLAG-TGF- β 1.

This example describes the construction and expression of two biologically functional TGF- β fusion proteins, each comprising the FLAG epitope tag near the
15 N-terminus of the mature TGF- β protein. The disclosed epitope-tagged TGF- β differs fundamentally from earlier attempts to tag TGF- β in that the inventors have succeeded in expressing epitope-tagged biologically active TGF- β in a mammalian host.

Expression of the FLAG-TGF- β fusions described in this example is driven
20 by the human elongation factor 1-alpha (EF1- α) gene promoter, a strong promoter capable of driving expression in virtually any type of cell. Two FLAG-tagged TGF- β 1 constructs, differing only in the location of insertion of the FLAG tag, are described. In one, the FLAG tag is inserted immediately following the cleavage site (N-terminal in the mature, processed TGF- β 1 molecule). In the second construct,
25 FLAG is inserted between amino acids 11 and 12 of the mature TGF- β 1 molecule. Each plasmid has successfully been transfected into mammalian cells, and the FLAG-TGF- β 1 protein product is detectable in cell culture supernatants using sandwich ELISA and western blots (with antibodies raised against either the FLAG epitope tag or TGF- β 1). Cells transfected with the fusions described in this example

secrete biologically active TGF- β 1 as measured by a number of different methods (see below).

Construction of FLAG-tagged TGF- β 1.

The FLAG epitope tag was introduced into the TGF- β 1 sequence using a 2-
5 step PCR mutagenesis technique. Flanking primers were:

(5'-primer) 5'-ggagagatctggtaccgagatggcgct-3 (SEQ ID NO: 1); and

(3'-primer) 5'-ataagaattgcgccgcttaatcgatcccaagtgggcttg-3' (SEQ ID NO:
2).

Two sets of FLAG mutagenesis primers were used. For constructing the N-
10 terminally tagged-TGF- β 1, the following primers were employed:

(5'-primer) 5'-gactacaaggatgacgacgacaaggccctggataccaactactgttc-3' (SEQ
ID NO: 3); and

(3'-primer) 5'-ctgtcgtcgtcatcctttagtctcggcggtgccgggagctgtg-3' (SEQ ID
NO: 4).

15 For the insertion of FLAG between amino acids 11 and 12 of the mature
TGF- β 1 peptide, the following mutagenesis primers were used:

(5'-primer) 5'-gactacaaggatgacgacgacaggagaagaactgtcgtgcggc-3' (SEQ ID
NO: 5);

(3'-primer) 5'-ctgtcgtcgtcatcctttagtctcggcggtgccgggagctgtg-3' (SEQ ID
20 NO: 6).

The PCR template was full length active porcine TGF- β 1, where two
cysteine residues at 223 and 225 in the LAP portion of the molecule have been
mutated to serines, disrupting two disulfide bonds that form between LAP and TGF-
25 β 1 (and which keep TGF- β 1 in a latent form), and where the second residue has
been changed from a proline to an alanine to improve translation. The final PCR
products had 5'-Bgl II and 3'-Not I restriction sites. The PCR products were
purified away from the primers using a QIAquick Gel Extraction kit (QIAGEN,
Valencia, CA) after separation on an agarose gel. Purified PCR products were
30 digested overnight and purified using a QIAGEN QIAquick PCR Purification kit
prior to ligation into BamHI, NotI digested, gel purified pEBB vector (Mayer *et al.*,

Curr. Biol., 5:296-305, 1995; provided by Colin Duckett, NIH). This vector contained the human elongation factor-1 α promoter, which drives expression of the cloned cDNA.

Transfection of Cos1 cells.

- 5 Cos1 cells were transiently transfected with each of the following: (1) pEBB vector alone; (2) WT (active) TGF- β 1 in pEBB (WT); (3) N-terminally FLAG-tagged TGF- β 1 (NFLAG β); and (4) TGF- β 1 with FLAG inserted between amino acids 11 and 12 of the mature TGF- β 1 peptide (11/12FLAG- β). Two micrograms of plasmid were transfected into cos1 cells in serum-free DMEM medium (Life Technologies, Rockville, MD) containing ITS+ supplement (Collaborative Biomedical Products, Bedford, MA) in 10 cm dishes using the Eugene 6 reagent and manufacturer's protocol (Roche, Indianapolis, IN). The total volume of media in each 10 cm dish was 5 mL. Supernatants were collected after 48 hours, and any contaminating cells were removed by brief centrifugation before storing
- 10
- 15 supernatants at -80° C.

Sandwich ELISA for active TGF- β 1.

Levels of active TGF- β 1 were quantified using a Quantikine TGF- β 1 sandwich ELISA kit from R&D Systems (Minneapolis, MN) following the instructions provided by the manufacturer.

- 20 *Antibodies and western blots.*

- All western blots were performed using Novex Tris-Glycine gels and the Novex blotting system (Novex, San Diego, CA). Protein was transferred onto nitrocellulose membranes, blocked with blocking buffer (TBST + 4% BSA) and incubated with the indicated antibodies in blocking buffer overnight at 4° C.
- 25 Secondary antibodies were conjugated to horseradish peroxidase (HRP) and purchased from Jackson Immunobiology. TGF- β 1 protein levels were measured using a rabbit polyclonal antibody (Cat. # G1221) purchased from Promega (Madison, WI). The anti-FLAG monoclonal antibody (M2) was purchased from Upstate Biotechnology (Waltham, MA). The anti-smad2 antibody was purchased
- 30 from Zymed (Cat # 51-1300, San Francisco, CA); and the anti-phospho-smad2 antibody was purchased from Upstate Biotechnology (Cat # 06-829).

Design of FLAG-tagged TGF- β .

The FLAG tag was inserted into two positions (FIG 2B) in different fusion constructs. In the first, FLAG was inserted immediately following the proteolytic cleavage site, which would make the tag N-terminal in the mature, processed TGF- β 1 molecule. In the second construct, FLAG was inserted between amino acids 11 and 12 of the mature TGF- β 1 peptide. This site was selected because, in one TGF- β family member, TGF- β 4, there is an insertion of an additional two amino acids - a phenomenon that does not affect biological activity of the native molecule.

FLAG-tagged TGF- β 1 is processed and secreted

10 Sandwich ELISA (SELISA) revealed that cells transfected with un-tagged TGF- β 1 (WT) secreted large amounts of the ligand into the medium (FIG 4A). Less ligand was secreted in the case of the NFLAG construct although the levels (5 ng/ml) are well in excess of the minimum concentration of ligand needed to inhibit cell growth (10-15 μ g/mL). In the case of the 11/12FLAG construct, SELISA was
15 unable to detect secreted TGF- β 1. It is possible that insertion of the FLAG sequence delays or impairs the production and/or secretion of TGF- β 1. However, we cannot rule out differences in transfection efficiencies at this time. Western blot analysis revealed that the un-tagged TGF- β 1 (WT) was indeed expressed to higher levels than were the FLAG-tagged constructs (FIG 4B). However, significant levels of
20 both the NFLAG and 11/12FLAG proteins were secreted into the medium as revealed with both the anti-TGF- β 1 antibody (panel A) and the anti-FLAG antibody (panel B). Consistent with the SELISA findings, levels of NFLAG- β 1 were higher than those of 11/12FLAG- β 1. Interestingly, the accumulation of the lower molecular weight, mature cleaved TGF- β 1 was lower in the case of cells transfected
25 with NFLAG- β 1 than in cells transfected with the 11/12FLAG- β 1 or WT constructs. It is possible that insertion of the FLAG sequence, which carries a strong net negative charge, interferes with cleavage of the pro-protein after the two positively charged arginine residues.

Example 2 Biological activity of FLAG-TGF- β 1 and 11/12FLAG-TGF- β 1.

This example provides assays that can be used to test the biological activity of TGF- β fusion molecules. In particular, this example demonstrates that representative functionalized TGF- β fusion proteins have TGF- β biological activity at least as great as native TGF- β , in spite of the addition of the FLAG epitope tag. Biological activity is demonstrated by two independent methods: (1) growth inhibition of CCL64 cells; and (2) phosphorylation of smad 2 in NMuMG cells.

Smad2 phosphorylation studies.

NmuMG cells were plated in 60-cm dishes, incubated in medium containing 0.5% FBS for three hours. COS cell supernatants from cells transfected with various TGF- β 1 plasmids were added to these cells (after first washing extensively with PBS). The cells were then incubated with the supernatants for 30 minutes. Cells were washed with PBS and scraped off the dishes. Cell pellets were dissolved in NP-40 lysis buffer containing 50 mM Tris-Cl pH 7.4, 150 mM NaCl, 50 mM NaF, 0.5% NP-40, 1 mM DTT, 1 mM sodium orthovanadate plus proteinase inhibitors (proteinase inhibitor cocktail tablets, one tablet per 10 mL buffer- Boehringer-Mannheim, Cat # 1836153, Indianapolis, IN). Thirty micrograms of total protein were applied to each lane of a 10% Novex Tris-Glycine gel.

Growth inhibition assays.

CCL64 mink lung epithelial cells were plated onto 6-well plates. Various dilutions of recombinant human TGF- β 1 standard or transfected Cos cell supernatants were added to the wells. CCL64 cells were pulsed for two hours with ^3H -thymidine, after 48 hours of incubation with TGF- β standard or conditioned medium. Cells were trypsinized and harvested using a cell harvester and [^3H] counts measured as described previously.

FLAG-tagged TGF- β 1 retains full biological activity

Supernatants from transfected COS cells were evaluated for biological activity by the ability to inhibit growth of and mink lung epithelial cell line, CCL64. Addition of recombinant human TGF- β 1 to these cells revealed that maximal inhibition was observed at about 35 $\mu\text{g/mL}$ (~ 97 $\mu\text{g/mL}$ for WT); 50% inhibition was observed at about 2.4 $\mu\text{g/mL}$ (~ 19 $\mu\text{g/mL}$ for WT). Comparison of the

supernatants from COS cells transfected with WT (un-tagged), NFLAG and 11/12 FLAG revealed that the FLAG-tagged ligands were at least as efficient as the wild type molecule in inhibiting cell growth (FIG 5).

Exposure of cells to TGF- β 1 leads to rapid and transient phosphorylation of smad 2 and smad 3. As a biochemical assay to measure TGF- β 1 activity, we looked at levels of smad 2 phosphorylation. NMuMG cells were treated with supernatants from transfected COS cells. After 30 minutes, the cells were washed extensively with PBS, scraped off the plates and the cell pellets were lysed in NP-40 lysis buffer. Smad 2 phosphorylation and total smad 2 level were examined by western blot analysis (FIG 4B, panel C). NMuMG cells incubated with supernatants from untransfected COS cells or from Cos cells transfected with vector alone (pEBB) showed no elevation in phosphorylated smad 2. However, cells incubated with supernatants from COS cells transfected with WT-TGF- β 1, NFLAG- β 1 and 11/12FLAG- β 1 all showed significant elevations in the amount of smad 2 phosphorylated (FIG 4B, panel C). Comparison of total smad 2 levels showed no significant differences between the samples. Thus, using two independent assays, one biological and the second biochemical, it is herein demonstrated that insertion of the FLAG epitope tag does not lead to significant loss of activity.

Example 3 Use of FLAG-TGF- β 1 and 11/12FLAG-TGF- β 1 in experimental systems.

This example demonstrates that the functionalizing peptide portion of the subject fusion proteins is functional in biological and test systems. FLAG-TGF- β 1 has been successfully used in a number of different assays. FLAG-TGF- β 1 was used to (1) detect expression of tagged ligand in transfected cells; (2) detect cell surface expression of TGF- β receptor complexes by flow cytometry; and (3) measure cell surface levels of receptor complexes in a non-radioactive cross-linking assay. The tagged ligand is a safer, faster and cheaper alternative to the use of [125 I] radiolabeled TGF- β 1 ligand and expands the repertoire of techniques that can be used to look at TGF- β receptor expression levels.

A. Flow cytometry: Detection of receptor molecule T β RI using FLAG-TGF- β 1

Cells were incubated with FLAG-TGF- β 1 in FACS buffer. Next, cells were incubated with a FITC-conjugated goat anti-mouse secondary antibody. This approach demonstrated that cell surface expression of TGF- β type II/type I receptor
5 complexes was detectable using the subject fusion proteins.

B. Immunocytochemistry: Detection of FLAG-tagged TGF- β 1

COS cells were transiently transfected with FLAG-tagged TGF- β 1 for 24 hours. Cells were then washed extensively with PBS and then fixed with buffered paraformaldehyde. Anti-TGF- β 1 or anti-FLAG monoclonal antibody was used,
10 followed by a secondary goat anti-mouse antibody conjugated to FITC, to detect intracellular expression FLAG- TGF- β 1 (FIG 4A, panels A and B respectively). The results clearly show that both epitope-tagged TGF- β fusion molecules easily can be detected in over-expressing cells.

C. Quantification of TGF- β receptor cell surface expression.

15 Cross-linking studies aimed at detecting cell surface expression of TGF- β receptors typically employ [125 I]-labeled TGF- β 1. As a safer and more economical alternative, FLAG-TGF- β 1 was examined to determine if it could be used in cross-linking assays. Cells were incubated with FLAG-TGF- β 1 in either the absence or presence of an excess of un-tagged TGF- β 1 ligand and cross-linking was performed
20 as previously described using DSS (Cheifetz *et al.*, *J. Bio. Chem.*, 261:9972-9978, 1986). Clarified cell lysates were first immunoprecipitated with a rabbit polyclonal antibody against the TGF β type II receptor (T β RII) (C-16, Santa Cruz). After extensive washing of the immune complexes, the samples were resuspended in 1X Laemmli buffer and resolved on a Novex 10% Tris-glycine gel. The blot was then
25 probed with an anti-FLAG monoclonal antibody. Both FLAG-TGF- β 1 ligands were effective in detecting cell surface TGF- β type I and type II receptor complexes. Detection of these complexes was abrogated by the addition of a 10-fold molar excess of un-tagged ligand. Film exposure times were in the range of two to four seconds, making this method of detection much faster than methods employing
30 radiolabeled ligand, which typically takes up to two weeks to develop.

Example 4 Additional Fusion Constructs

Additional functionalized TGF- β fusion proteins have been constructed using different TGF- β isoforms (*e.g.*, TGF- β 2 and TGF- β 3) and different functionalizing peptide portions (*e.g.*, haemagglutinin (HA) and Green Fluorescent protein (GFP)), as well as placing the tag in slightly different positions than at the N-terminus or between residues 11 and 12. In particular examples of such fusion constructs, the tag has been introduced between amino acid 5 and 6 of mature TGF- β 1, or optionally at the front of the mature protein with a short amino acid sequence (*e.g.*, ten or fewer, for instance five amino acids) from the mature TGF- β family protein duplicated before the functionalizing peptide.

This example provides descriptions of some of the preparation and/or characterization of additional fusion constructs.

15 Methods and Materials:

Construction of Fusion Protein Constructs: For construction of the plasmid containing the N+5FLAG TGF- β 1 (N+5FLAG- β 1) construct (SEQ ID NO: 16), the following mutagenesis primers were used:

(forward) 5'-gccctggataccaacgactacaaggatgacgacgacaaggccctggataccaactactgcttcag-
ctccacgg-3' (SEQ ID NO: 18);
(reverse) 5'-cttgtcgtcgtcatccttgtatgctgtatccagggctcggcggtggtgccgggagctgtgcagg-
gctgggc-3' (SEQ ID NO: 19).

For construction of the plasmid containing the N+5HA-TGF- β 1 (N+5HA- β 1) construct (SEQ ID NO: 20), the following primer pairs were used:
(forward: 5'-gccctggataccaacagctaccatacagctgccagactacgcatctctggccctggataccaact-
actgcttcagctccacggagaagaactgctgcgtgcggcag-3' (SEQ ID NO: 22);
(reverse) 5'-cagagatgcgtagtctggcacgtcgtatgggtagctgttggtatccagggctcggcggtgccggga-
gctgtgc-3' (SEQ ID NO: 23).

The same pair of flanking primers as those employed for construction of the N-FLAG- β 1 construct (Example 1) was used. The gel purified PCR products were cloned into the mammalian expression vector pEF6-V5/His-TOPO (InVitrogen,

Carlsbad, CA). The presence of a stop codon in the porcine cDNA ensured that the c-terminal V5 and 6x His tags in the vector were not translated.

Transfection of Cos1 cells. Cos1 cells were transiently transfected with pEBB or pEF6-V5/His-TOPO vector alone; WT (active) TGF- β 1 in pEBB or PEF6-V5/His-TOPO (WT); N-terminally FLAG-tagged TGF- β 1 (NFLAG β); N+5FLAG-TGF β 1; or N+5 HA-TGF β 1. Two micrograms of plasmid were transfected into Cos1 cells in serum-free DMEM medium containing ITS+ supplement (Collaborative Biomedical Products) in 10 cm dishes using the Eugene 6 reagent and manufacturer's protocol (Roche). The total volume of media in each 10 cm dish was 5 mL. Supernatants were collected after 48 hours, and any contaminating cells were removed by brief centrifugation before storing supernatants at -80° C.

Sandwich ELISA for active epitope-tagged TGF- β 1. Levels of FLAG- and HA-tagged TGF- β 1 were measured using a sandwich ELISA. A flat 96-well plate (Falcon) was used; each well was coated with 50 μ l of the first antibody (mouse-anti-FLAG M2, or mouse-anti-HA, each at 20 μ g/ml in PBS). Uncoated surfaces of the wells were blocked by addition of 250 μ l per well of blocking buffer (PBS + 3% BSA) for 30 minutes. The conditioned media were used in serial dilutions, starting at 1X. As a secondary antibody we used 50 μ l of anti-TGF- β 1-antibody-conjugate (R&D SYSTEMS, Quantikine ELISA kit). Equal amounts of stabilized hydrogen peroxide and stabilized chromogen (tetramethylbenzidine) (both R&D SYSTEMS) were added. After 20 minutes of incubation stop solution (R&D SYSTEMS) was added. The optical density of each well was determined using a microplate reader, set to 450 nm.

Antibodies and western blots. All western blots were performed using Novex Tris-Glycine gels and the Novex blotting system. Protein was transferred onto nitrocellulose membranes, blocked with blocking buffer (TBST + 4% BSA) and incubated with the indicated antibodies in blocking buffer overnight at 4° C. Secondary antibodies were conjugated to HRP and purchased from Jackson Immunolabs. After 5 washes with deionized water, blots were incubated for 1 hour with secondary antibody diluted 1:10,000 in blocking buffer. Blots were then washed four times for 5 minutes each in deionized water, followed by a 15 minutes

wash in TBS + 0.1% Tween 20 and a final series of four washes in deionized water (5 minutes per wash). Membranes were blotted to remove excess water and incubated with substrate solution (SuperSignal West Pico, Pierce) according to the manufacturer's instructions. After blotting the membranes to remove excess solution, the blots were exposed to x-ray film (Bio-Max MR, Kodak). TGF- β 1 protein levels were measured using a rabbit polyclonal antibody (Cat# G1221) purchased from Promega. The anti-FLAG monoclonal antibody (M2) was purchased from Upstate Biotechnology. The anti-smad2 antibody was purchased from Zymed (Cat # 51-1300); and the anti-phospho-smad2 antibody was purchased from Upstate Biotechnology (Cat # 06-829).

Smad2 phosphorylation studies. Mv1Lu cells were plated onto 6-well plates, in DMEM medium containing 0.2% FBS. Twenty-four hours later, conditioned media from Cos1 cells transfected with various TGF- β 1 plasmids were diluted twenty-fold and added to these cells (after first washing extensively with PBS). The cells were then incubated with the conditioned media for 30 minutes. Cells were washed with PBS and harvested. Cell pellets were dissolved in NP-40 lysis buffer containing 50 mM Tris-Cl pH 7.4, 150 mM NaCl, 50 mM NaF, 0.5% NP-40, 1 mM DTT, 1 mM sodium orthovanadate plus proteinase inhibitors (proteinase inhibitor cocktail tablets, 1 tablet per 10 mL buffer- Boehringer-Mannheim, Cat # 1836153). Thirty micrograms of total protein were applied to each lane of a 10% Novex Tris-Glycine gel.

Growth inhibition assays. Mv1Lu mink lung epithelial cells (5×10^5 cells per well) were plated onto 96-well plates in DMEM containing 0.2% FBS. Various dilutions of recombinant human TGF- β 1 standard or transfected Cos cell supernatants were added to the wells. After 24 hours, cells were pulsed with 1 μ Ci of [3 H]thymidine for 4 hours. Cells were trypsinized and harvested using a Packard Bioscience cell harvester and [3 H]-thymidine counts measured by liquid scintillation spectrometry.

Flow cytometry. Mv1Lu cells were harvested using CellStripper (Manufacturer and location?), a non-enzymatic dissociation solution. Cells were counted and 3×10^6 cells were transferred to 5 mL FACS tubes. Cells were washed

with PBS and fixed with 4% buffered paraformaldehyde on ice, for 5 minutes. The paraformaldehyde (8%) was added dropwise to cells in an equal volume of PBS with gentle vortexing to avoid clumping of cells. Cell pellets were then washed with 4 mL of cold PBS. Cells were permeabilized by the addition of 1 mL of -20C

5 methanol with vortexing and incubated on ice for 2 minutes. Cells were washed with PBS and then treated for 5 minutes at room temperature with 50 mM glycine in PBS (to quench auto-fluorescence). After washing with PBS, cells were pelleted and 200 μ l of blocking buffer plus 20 μ l of conditioned medium containing N+5FLAG- or N+5HA-TGF- β 1 ligand were added. Cells were incubated with

10 ligand for 2.5 hours at 4° C. Following incubation with ligand, cells were washed two times with ice-cold PBS and once with blocking buffer. Cells were then incubated with blocking buffer for 30 minutes at room temperature prior to addition of 200 μ l of blocking buffer containing primary antibody (anti-FLAG, anti-HA monoclonal or anti-KLH isotype control) diluted 1:1000 (approximately 0.4 μ g in

15 200 μ L). Incubation with primary antibody was carried out at 4° C overnight. The following day, samples were washed twice with ice-cold PBS and once with blocking buffer. Samples were then incubated for one hour with TRITC-conjugated goat anti-mouse (Jackson Immunolabs) at 1:100 in blocking buffer. After washing twice with PBS and once with FACS buffer (PBS + 1.0% heat-inactivated FBS),

20 samples were resuspended in 400 μ l of FACS buffer and analyzed by flow cytometry using a FACSCalibur (Beckton Dickinson).

Confocal immunofluorescence microscopy. To confirm intracellular detection Cos1 cells were split on coverslips and transfected with either WT TGF- β 1, FLAG- or HA-tagged TGF- β 1 or WT-TGF- β 1 and one of the tagged TGF- β 1

25 plasmids. The next day the cells were fixed with 3.5% paraformaldehyde in PBS and permeabilized with methanol (-20°C). Glycine (50 mM in PBS) was used to quench auto-fluorescence (5 minutes at room temperature). Non-specific binding sites were blocked with 10% normal goat serum in PBS and the cells were then incubated overnight with the primary antibody; mouse-anti-FLAG, mouse-anti-HA

30 or rabbit-anti-TGF- β 1. As secondary antibodies, FITC-conjugated goat-anti-rabbit and TRITC-conjugated goat-anti-mouse were used. Coverslips were mounted on

slides, using mounting medium with DAPI (Vector Labs) and analyzed using confocal immunofluorescence microscopy.

Results:

5 *Design of FLAG-tagged TGF- β .*

Initially, we chose to insert the FLAG tag immediately following the proteolytic cleavage site, which would make the tag N-terminal in the mature, processed molecule (FIG 6A).

FLAG-tagged TGF- β 1 is processed and secreted

10 Immunoblot analyses revealed that cells transfected with un-tagged TGF- β 1 (WT) secreted large amounts of the ligand into the medium (FIG 6B). Less ligand was secreted in the case of the NFLAG, although the levels, as determined by TGF- β sandwich ELISA (5 ng/ml) were well in excess of the minimum concentration of ligand (78 pg/mL) needed to maximally inhibit cell growth in these experiments
15 according to a standard curve using recombinant human TGF- β 1. Indeed, this N-terminally tagged TGF- β 1 ligand is biologically active as it induces significant phosphorylation of smad 2, a TGF- β signaling intermediate (FIG 6B). Levels of total smad 2 were equivalent in all treatments.

There was less accumulation of the lower molecular weight, mature cleaved
20 TGF- β 1 in the case of the FLAG-tagged molecule, relative to the un-tagged ligand. The cleavage site that separates LAP from the mature TGF- β peptide is preceded by a region of positively charged amino acids. Without meaning to be limited to one possible explanation, it is hypothesized that insertion of the FLAG epitope sequence, which carries a strong net negative charge, interferes with efficient cleavage of the
25 pro-protein.

The reduced cleavage efficiency in the NFLAG protein was further analyzed by placing the FLAG tag further downstream of the cleavage site. Additional constructs were generated in which the FLAG or HA tag was inserted after the first five amino acids downstream of the cleavage site (N+5FLAG and N+5HA
30 constructs). In addition, immediately following the tags, the first five amino acids of the mature peptide were re-iterated, followed by the remaining sequence of the

mature peptide (FIG 6C). These re-designed molecules appear to exhibit cleavage efficiencies comparable to that of the un-tagged molecule (FIG 6D). In addition, the levels of secreted TGF- β 1 appear equivalent for tagged and un-tagged molecules.

FLAG- and HA-tagged TGF β 1 retain biological activity

5 Comparison of the supernatants from Cos1 cells transfected with WT (un-tagged), N+5FLAG- and N+5HA-TGF- β 1 revealed that the FLAG- and HA-tagged ligands were as efficient as the wild type molecule in inhibiting cell growth (FIG 7A). Inhibition of growth was reversed by inclusion of a TGF- β neutralizing antibody, indicating a specific effect due to active TGF- β . No inhibition of growth
10 was seen for media from Cos1 cells transfected with empty vector, suggesting that any endogenous TGF- β from Cos1 cells must be below the limits of detection in our assays.

 TGF- β 1 treatment leads to rapid and transient phosphorylation of smad 2 and smad 3. As a biochemical assay to measure TGF- β 1 activity, we looked at levels of
15 smad 2 phosphorylation (FIG 7B). Mv1Lu cells incubated with conditioned medium from Cos1 cells transfected with vector alone showed no elevation in phosphorylated smad 2. However, cells incubated with media containing WT-TGF- β 1, N+5FLAG- β 1 and N+5HA-TGF- β 1 showed significant elevations in the amount of phosphorylated smad 2. The accumulation of phospho-smad 2 was abrogated by
20 the presence of TGF- β neutralizing antibody, indicating that phosphorylation was indeed specific to TGF- β treatment. Comparison of total smad 2 protein levels showed no significant differences between the samples. Thus, using two independent assays, one biological and the second biochemical, we have shown that insertion of the FLAG and HA epitope tags does not lead to loss of activity.

25 *Specific detection of FLAG- and HA-tagged TGF- β 1 by sandwich ELISA (SELISA).*

 A particularly important application of the herein provided, newly designed TGF- β ligands is their use as therapeutics, with the ability to track the distribution of the administered cytokine with tag-specific reagents. To exploit this new use, a sandwich ELISA assay was developed to specifically detect epitope-tagged TGF- β 1.
30 Epitope-tagged ligand was captured by antibody immobilized onto the wells of a 96

well plate. Detection was facilitated by the use of the TGF- β 1 conjugate, a polyclonal antibody against TGF- β 1 conjugated to horse radish peroxidase (HRP) supplied in the human TGF- β 1 Quantikine ELISA kit (R&D Systems). This assay was able to specifically detect FLAG- or HA-tagged TGF- β 1 ligand in conditioned medium from transfected Cos-1 cells (FIG 7C).

Detection of the FLAG-tagged molecule was possible at a dilution of 1:50, whereas specific detection of the HA-tagged ligand was achievable up to a 1:10 dilution. The inability to detect the HA-tagged ligand at lower concentrations may have been a result of lower levels of ligand in conditioned medium from this particular batch of conditioned medium. This again could be explained by variable transfection efficiencies between experiments. Alternatively, it is possible that the anti-HA monoclonal used here has a lower binding efficiency to plastic.

FLAG-tagged TGF- β 1 can be detected by immunofluorescence confocal microscopy

The utility of the tagged ligands was next tested in an immunocytochemical application. Cos-1 cells grown on glass coverslips were transfected with WT TGF- β 1 only, the WT- and N+5FLAG-tagged TGF- β 1 or both the WT- and HA-tagged TGF- β 1. A FITC-conjugated goat anti-rabbit antibody was used to bind and detect TGF- β 1 and a TRITC-conjugated goat anti-mouse antibody was used to bind and detect either the FLAG or the HA-epitope. Antibodies directed against either of the epitope tags do not detect the untagged TGF- β 1 molecule; whereas, the anti-TGF- β 1 antibody is able to clearly detect both the untagged as well as the tagged molecules. Antibodies directed against either FLAG or HA were able to detect the appropriate tagged TGF- β 1. Both untagged and epitope-tagged molecules show a high degree of co-localization.

Use of FLAG- and HA- tagged TGF- β 1 to detect cell surface and cytoplasmic expression of T β RII by flow cytometry

In addition to their potential as therapeutic reagents, the herein provided tagged ligands can serve as valuable diagnostic reagents for probing the pattern and level of TGF- β receptor expression *in vivo*. To demonstrate this, the FLAG- and

HA-tagged TGF- β 1 fusion proteins were used to detecting cell surface and cytoplasmic expression of T β RII by flow cytometry.

Mv1lu cells were harvested using a non-enzymatic cell dissociation solution to avoid proteolysis of surface receptor complexes. Cells were fixed and
5 permeabilized to examine total levels of TGF- β receptors (surface + intracellular), or fixed only (without permeabilization) to permit specific detection of surface receptor only. Cells were incubated with blocking buffer (10% goat serum in PBS) followed by incubation with blocking buffer containing a 1:100 dilution of conditioned medium containing epitope-tagged ligand. After washing, the primary antibody
10 directed against the epitope tag was added and detection was facilitated using a TRITC-conjugated goat anti-mouse IgG. Labeling with FLAG-TGF- β 1 results in a shifted peak relative to isotype control (mouse anti-KLH) indicating the presence TGF- β receptor complexes (surface + intracellular) and a smaller shift in the case of non-permeabilized cells (cell surface receptor). Similar results were observed for
15 the HA-TGF- β 1 ligand. A greater shift in fluorescence intensity was consistently observed for fixed and permeabilized cells, a result consistent with previous studies showing that the majority of TGF- β receptor molecules are localized intracellularly and not on the cell surface (Zwaagstra *et al.*, *Exp. Cell Res.*, 121-134, 2000).

Examples 3 and 4 demonstrates that several TGF- β fusions successfully used
20 in a number of different assays, including:

- (1) detection of expression of tagged ligand in transfected cells;
- (2) detection of expression of TGF- β receptor complexes by flow cytometry;
- and
- (3) measurement of levels of tagged ligand in tissue culture supernatants.

25 It is believed that the flow cytometry assay described here represents a safer, faster, and cheaper alternative to the use of [125 I] radiolabeled TGF- β 1 ligand, and expands the repertoire of techniques that can be used to look at TGF- β receptor expression levels. These tagged ligands will circumvent the complications due to
30 cross reactivity of certain isoform-specific TGF- β antibodies. Finally, reagents will

prove important in developing new transgenic models in which epitope-tagged TGF- β isoforms can be expressed in a spatially and/or temporally restricted manner.

Example 5 Further TGF- β Fusions

Sub 5 Additional functionalized TGF- β fusion proteins have been constructed in a manner essentially similar to the methods described in Examples 3 and 4. Specific examples of additional functionalized fusions include:

- 10 murine N+5FLAG TGF- β 2 (MN5FLAGb2; SEQ ID NO: 24 and 25);
murine N+5HA TGF- β 2 (MN5HAb2; SEQ ID NO: 26 and 27);
murine N+5FLAG TGF- β 3 (MN5FLAGb3; SEQ ID NO: 28 and 29);
murine N+5HA TGF- β 3 (MN5HAb3; SEQ ID NO: 30 and 31);
porcine active N+5FLAG TGF- β 1 (actN5FLAGb1; SEQ ID NO: 32 and 33);
porcine latent N+5FLAG TGF- β 1 (latN5FLAGb1; SEQ ID NO: 34 and 35)
(made latent by mutations at positions 682 and 688 of SEQ ID NO: 34);
15 porcine active N+5FLAG TGF- β 1 (actN5HAb1; SEQ ID NO: 36 and 37);
and
porcine latent N+5FLAG TGF- β 1 (latN5HAb1; SEQ ID NO: 38 and 39)
(made latent by mutations at positions 678 and 684 of SEQ ID NO: 38).

Sub 5 Additional characteristics of each of these fusions, as well as N+5FLAG-TGF- β 1 (SEQ ID NOS: 18 and 19) and N+5HA-TGF- β 1 (SEQ ID NO: 22-23) fusions, are provided in Table 2.

Table 2

Fusion name and SEQID NOs.	5'UTR	CDS	3'UTR	AA 1-5 of TGF- β	Epitope tag	Mature fusion
N+5FLAG-TGF- β 1 (NOs: 18 & 19) ¹	1-347 ²	348-1559	1560-1612	1182-1196	1197-1220	182-1559
N+5HA-TGF- β 1 (NOs: 22 & 23)	1-347	347-1571	1572-1624	1182-1196	1197-1232	1182-1571
N+5FLAG TGF- β 2 (NOs: 24 & 25)	N/A	1-1284	N/A	907-921	922-945	907-1284
N+5HA TGF- β 2 (NOs: 26 & 27)	1-7	8-1303	N/A	914-928	929-964	914-1303
N+5FLAG TGF- β 3 (NOs: 28 & 29)	N/A	1-1272	N/A	895-909	910-945	895-1272
N+5HA TGF- β 3 (NOs: 30 & 31)	N/A	1-1284	N/A	895-909	910-945	895-1284
N+5FLAG TGF- β 1 (NOs: 32 & 33)	1-10	11-1222	1223-1349	845-859	860-883	845-1222
N+5FLAG-TGF- β 1 (NOs: 34 & 35)	1-14	15-1226	1227-1253	849-863	864-887	849-1226
N+5FLAG TGF- β 1 (NOs: 36 & 376)	1-10	11-1234	1235-1361	845-859	860-895	845-1234
N+5FLAG TGF- β 1 (NOs: 38 & 39)	1-10	11-1234	1335-1361	845-859	860-895	845-1234

¹Refers to the nucleic acid sequence and amino acid sequence for the listed fusion.

²Residue positions correspond to the position in the nucleic acid sequence.

- 5 This disclosure provides methods for producing functionalized small molecules, including peptide-tagged TGF- β family proteins and other cytokines. The disclosure further provides such functionalized molecules, and methods of using these molecules in the clinical setting, as diagnostic and prognostic tools, therapy monitors, and so forth. It will be apparent that the precise details of the methods
- 10 described may be varied or modified without departing from the spirit of the described invention. We claim all such modifications and variations that fall within the scope and spirit of the claims below.

Sub
B6

FOR OFFICIAL USE ONLY